




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 233/60, 233/64, 417/06, 401/12, A61K 31/415		A1	(11) International Publication Number: WO 99/20611
			(43) International Publication Date: 29 April 1999 (29.04.99)
(21) International Application Number: PCT/GB98/03117		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 19 October 1998 (19.10.98)		Published With international search report.	
(30) Priority Data: 97402505.8 22 October 1997 (22.10.97) EP 97402502.5 22 October 1997 (22.10.97) EP			
(71) Applicants (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). ZENECA-PHARMA S.A. [FR/FR]; Le Galien, 1, rue des Chauffours, Boîte postale 127, F-95022 Cergy Cedex (FR).			
(72) Inventor; and (75) Inventor/Applicant (for US only): ARNOULD, Jean-Claude [FR/FR]; Centre de Recherches, Zone Industrielle La Pompe, Chemin de Vrilly, Boîte postale 1050, F-51689 Reims Cedex 2 (FR).			
(74) Agent: TINSLEY, Rachel, Maria; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).			

(54) Title: IMIDAZOLE DERIVATIVES AND THEIR USE AS FARNESYL PROTEIN TRANSFERASE INHIBITORS



(57) Abstract

The present invention relates to compound of formula (1), wherein Ar¹ represents (A), or (B), or (C); R¹² and R¹³ are independently hydrogen or C₁₋₄ alkyl; Ar² is phenyl or heteroaryl; p is 0 or 1; Ar³ is phenyl, pyridinyl, pyridazinyl, pyrimidyl or pyrazynyl the ring being substituted on ring carbon atoms by R² and -(CH₂)_nR³ and wherein Ar³ is attached to Ar¹C(R¹²)R¹³CH(Ar²)O- by a ring carbon atoms; R² is a group of formula (2), or R² represents a lactone of formula (3), the group of formula (2) or (3) having *L* or *D* configuration at the chiral alpha carbon in the corresponding free amino acid; n is 0, 1 or 2; R³ is phenyl or heteroaryl; and R⁵-R⁹, m and n are as defined in the specification; or a pharmaceutically acceptable salt, prodrug or solvate thereof. Processes for their preparation, their use as therapeutic agents and pharmaceutical compositions containing them. A particular use is in cancer therapy.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NR	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

IMIDAZOLE DERIVATIVES AND THEIR USE AS FARNESYL PROTEIN TRANSFERASE
INHIBITORS

This invention relates to compounds that inhibit farnesylation of mutant ras gene products through inhibition of the enzyme farnesyl-protein transferase (FPTase). The invention also relates to methods of manufacturing the compounds, pharmaceutical compositions and methods of treating diseases, especially cancer, which are mediated through farnesylation of ras.


Cancer is believed to involve alteration in expression or function of genes controlling cell growth and differentiation. Whilst not wishing to be bound by theoretical considerations the following text sets out the scientific background to ras in cancer. Ras genes are frequently mutated in tumours. Ras genes encode guanosine triphosphate (GTP) binding proteins which are believed to be involved in signal transduction, proliferation and malignant transformation. H-, K- and N-ras genes have been identified as mutant forms of ras (Barbacid M, Ann. Rev. Biochem. 1987, 56: 779-827). Post translational modification of ras protein is required for biological activity. Farnesylation of ras catalysed by FPTase is believed to be an essential step in ras processing. It occurs by transfer of the farnesyl group of farnesyl pyrophosphate (FPP) to a cysteine at the C-terminal tetrapeptide of ras in a structural motif called the CAAX box. After further post-translational modifications, including proteolytic cleavage at the cysteine residue of the CAAX box and methylation of the cysteine carboxyl, ras is able to attach to the cell membrane for relay of growth signals to the cell interior. In normal cells activated ras is believed to act in conjunction with growth factors to stimulate cell growth. In tumour cells it is believed that mutations in ras cause it to stimulate cell division even in the absence of growth factors (Travis J, Science 1993, 260: 1877-1878), possibly through being permanently in GTP activated form rather than cycled back to GDP inactivated form. Inhibition of farnesylation of mutant ras gene products will stop or reduce activation.

One class of known inhibitors of farnesyl transferase is based on farnesyl pyrophosphate analogues; see for example European patent application EP 534546 from Merck. Inhibitors of farnesyl transferase based on mimicry of the CAAX box have been reported. Reiss (1990) in Cell 62, 81-8 disclosed tetrapeptides such as CVIM (Cys-Val-Ile-Met). James (1993) in Science 260, 1937-1942 disclosed benzodiazepine based peptidomimetic compounds. Lerner (1995) in J. Biol. Chem. 270, 26802 and Eisai in

- 2 -


International Patent Application WO 95/25086 disclosed further peptidomimetic compounds based on Cys as the first residue. Bristol-Myers Squibb in European Patent Application EP 696593 disclosed farnesyl transferase inhibitors having a 4-sulfanylpyrrolidine residue in the first position.

- 5 According to one aspect of the present invention there is provided a compound of Formula (1):



Formula (1)

wherein Ar¹ represents:



R⁵ is hydrogen, C₁₋₄alkyl, phenylC₁₋₄alkyl;

- 10 R⁶ is hydrogen, C₁₋₄alkyl, hydroxyC₁₋₄alkyl, haloC₁₋₄alkyl, dihaloC₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, sulfanylC₁₋₄alkyl, aminoC₁₋₄alkyl, N-(C₁₋₄alkyl)aminoC₁₋₄alkyl, N,N-di(C₁₋₄alkyl)aminoC₁₋₄alkyl or phenylC₁₋₄alkyl; m is 0, 1 or 2;

R¹² and R¹³ are independently hydrogen or C₁₋₄ alkyl;

Ar² is phenyl or heteroaryl;


- 15 p is 0 or 1;

Ar³ is phenyl, pyridinyl, pyridazinyl, pyrimidyl or pyrazinyl, the ring being substituted on ring carbon atoms by R² and -(CH₂)_nR³ and wherein Ar³ is attached to

Ar¹C(R¹²)R¹³CH(Ar²)O- by a ring carbon atom;


R² is a group of the Formula (2):

- 3 -



Formula (2)

- wherein R^7 is hydrogen or C_{1-4} alkyl, R^8 is $-(CH_2)_q-R^{10}$ wherein q is 0-4 and R^{10} is C_{1-4} alkylsulfanyl, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, hydroxy, C_{1-4} alkoxy, carbamoyl, \underline{N} - C_{1-4} alkyl carbamoyl, $\underline{N,N}$ -(di C_{1-4} alkyl)carbamoyl, C_{1-4} alkyl, phenyl, thienyl, or
- 5 C_{1-4} alkanoylamino, R^9 is hydroxy, C_{1-6} alkoxy, C_{3-9} cycloalkyloxy, heterocyclyloxy, heterocyclyl C_{1-4} alkoxy or $-NH-SO_2-R^{11}$ wherein R^{11} represents, trifluoromethyl, C_{1-4} alkyl, phenyl, heteroaryl, aryl C_{1-4} alkyl or heteroaryl C_{1-4} alkyl;
- or R^2 represents a lactone of Formula (3)



Formula (3)

- 10 the group of Formula (2) or (3) having \underline{L} or \underline{D} configuration at the chiral alpha carbon in the corresponding free amino acid;
- n is 0, 1 or 2;
- R^3 is phenyl or heteroaryl;
- phenyl and heteroaryl rings in R^3 , R^5 , R^6 , R^9 , R^{11} and Ar^2 are independently optionally
- 15 substituted on ring carbon atoms in by up to three substituents selected from C_{1-4} alkyl, halogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, amino, C_{1-4} alkylamino, di(C_{1-4} alkyl)amino, C_{1-4} alkanoylamino, nitro, cyano, carboxy, thiol, C_{1-4} alkylsulfanyl, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkanesulphonamido, \underline{N} -(C_{1-4} alkylsulphonyl)- \underline{N} - C_{1-4} alkylamino, aminosulfonyl, \underline{N} -(C_{1-4} alkyl)aminosulfonyl,
- 20 $\underline{N,N}$ -di(C_{1-4} alkyl)aminosulfonyl, carbamoyl, \underline{N} -(C_{1-4} alkyl)carbamoyl, $\underline{N,N}$ -(di C_{1-4} alkyl)carbamoyl, carbamoyl C_{1-4} alkyl, \underline{N} -(C_{1-4} alkyl)carbamoyl C_{1-4} alkyl, $\underline{N,N}$ -(di C_{1-4} alkyl)carbamoyl C_{1-4} alkyl, hydroxy C_{1-4} alkyl and C_{1-4} alkoxy C_{1-4} alkyl and on ring NH groups (replacing hydrogen) by C_{1-4} alkyl, C_{1-4} alkanoyl, C_{1-4} alkylsulfonyl, halo C_{1-4} alkyl, difluoromethyl or trifluoromethyl;
- 25 or a pharmaceutically-acceptable salt, prodrug or solvate thereof.

In this specification the generic term "alkyl" includes both straight-chain and branched-chain alkyl groups. However references to individual alkyl groups such as "propyl" are specific for the straight-chain version only and references to individual branched-chain alkyl groups such as "isopropyl" are specific for the branched-chain version only. An
 5 analogous convention applies to other generic terms.

It is to be understood that, insofar as certain of the compounds of Formula (1) defined above may exist in optically active or racemic forms by virtue of one or more asymmetric carbon atoms, the invention includes in its definition any such optically active or racemic form which possesses the property of inhibiting FTPase. The synthesis of optically
 10 active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. Similarly, inhibitory properties against FTPase may be evaluated using the standard laboratory techniques referred to hereinafter.

The term "heterocyclyl" refers to a 5- or 6-membered monocyclic ring containing 1 to
 15 3 heteroatoms selected from nitrogen, oxygen and sulfur.

The term "heteroaryl" refers to a 5-10 membered monocyclic heteroaryl ring containing upto 3 heteroatoms selected from nitrogen, oxygen and sulphur.

The term " halogen " refers to fluorine, chlorine, bromine and iodine. The term "carbamoyl" refers to -C(O)NH₂. The term " BOC " refers to tert-butoxycarbonyl.

20 Examples of **C₁₋₄alkyl** include methyl, ethyl, propyl, isopropyl, *sec*-butyl and *tert*-butyl; examples of **C₁₋₄alkoxy** include methoxy, ethoxy and propoxy; examples of **C₁₋₄alkanoyl** include formyl, acetyl and propionyl; examples of **C₁₋₄alkanoyloxy** include acetyloxy and propionyloxy; examples of **C₁₋₄alkylamino** include methylamino, ethylamino, propylamino, isopropylamino, *sec*-butylamino and *tert*-butylamino; examples of **di-(C₁₋₄alkyl)amino** include di-methylamino, di-ethylamino and N-ethyl-N-methylamino; examples
 25 of **C₁₋₄alkanoylamino** include acetamido and propionylamino; examples of **C₁₋₄alkoxycarbonyl** include methoxycarbonyl, ethoxycarbonyl and propoxycarbonyl; examples of **C₁₋₄alkylsulfanyl** include methylsulfanyl, ethylsulfanyl, propylsulfanyl, isopropylsulfanyl, *sec*-butylsulfanyl and *tert*-butylsulfanyl; examples of **C₁₋₄alkylsulfinyl**
 30 include methylsulfinyl, ethylsulfinyl, propylsulfinyl, isopropylsulfinyl, *sec*-butylsulfinyl and

tert-butylsulfinyl; examples of **C₁₋₄alkylsulfonyl** include methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, *sec*-butylsulfonyl and *tert*-butylsulfonyl; examples of **N-(C₁₋₄alkyl)carbamoyl** include N-methylcarbamoyl and N-ethylcarbamoyl; examples of **N,N-(diC₁₋₄alkyl)carbamoyl** include N,N-dimethylcarbamoyl and N-methyl-N-ethylcarbamoyl;

5 examples of **C₁₋₄alkanesulfonamido** include methanesulfonamido, ethanesulphonamido and propanesulfonamido; examples of **C₁₋₄alkylsulfonyl-N-C₁₋₄alkylamino** include methylsulfonyl-N-methylamino, ethylsulfonyl-N-methylamino and propylsulfonyl-N-methylamino; examples of fluoroC₁₋₄alkyl include fluoromethyl, 2-fluoroethyl and 3-fluoropropyl; examples of difluoroC₁₋₄alkyl include difluoromethyl, 2,2-difluoroethyl and

10 3,3-difluoropropyl; examples of **carbamoylC₁₋₄alkyl** include carbamoylmethyl, carbamoylethyl and carbamoylpropyl; examples of **N-(C₁₋₄alkyl)carbamoylC₁₋₄alkyl** include N-methyl-carbamoylmethyl and N-ethyl-carbamoylethyl; examples of **N,N-(diC₁₋₄alkyl)carbamoyl-C₁₋₄alkyl** include N,N-dimethylcarbamoylethyl and N-methyl-N-ethylcarbamoylethyl; examples of **hydroxyC₁₋₄alkyl** include hydroxymethyl, hydroxyethyl,

15 hydroxypropyl, 2-hydroxypropyl, 2-(hydroxymethyl)propyl and hydroxybutyl; examples of **C₁₋₄alkoxyC₁₋₄alkyl** include methoxyethyl, ethoxyethyl and methoxybutyl; examples of **sulfanylC₁₋₄alkyl** include sulfanylmethyl, sulfanylethyl, sulfanylpropyl; and examples of **N-(C₁₋₄alkyl)aminoC₁₋₄alkyl** include N-methyl-aminomethyl and N-ethyl-aminoethyl.

Examples of 5- or 6-membered heteroaryl ring systems include imidazole, triazole,

20 pyrazine, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole and thiophene.

Preferably the NH group in imidazole is unsubstituted or substituted by C₁₋₄alkyl.

Examples of heterocyclyl rings include pyrrolidinyl, morpholinyl, piperidinyl, dihydropyridinyl and dihydropyrimidinyl.

25 Preferred heteroatoms are N and S, especially N. In general, attachment of heterocyclic rings to other groups is via carbon atoms.

Examples of values for R⁸ in Formula (2) are side chains of lipophilic amino acids including such as for example methionine, phenylglycine, phenylalanine, serine, leucine, isoleucine or valine. L configuration in the corresponding free amino acid is preferred.

30 Examples of amino acid side chains are set out below.

Amino Acid	Side Chain
methionine	$-\text{CH}_2-\text{CH}_2-\text{S}-\text{CH}_3$
phenylglycine	Ph
phenylalanine	$-\text{CH}_2-\text{Ph}$
thienylalanine	$-\text{CH}_2-\text{thien-2-yl}$
serine	$-\text{CH}_2\text{OH}$ or a C_{1-4} alkyl (preferably methyl) ether thereof.
Leucine	$-\text{CH}_2-\text{CHMe}_2$
homoserine	$-\text{CH}_2-\text{CH}_2-\text{OH}$ or a C_{1-4} alkyl (preferably methyl) ether thereof
<u>N</u> -acetyl-lysine	$-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CO}-\text{CH}_3$

The lactone in Formula (3) can be formed from a group of Formula (2) when R^9 is OH to give a carboxyl and R^8 is $-\text{CH}_2-\text{CH}_2-\text{OH}$ where R^8 and R^9 together lose a water molecule to form part of a dihydrofuran-2-one heterocyclic ring.

Preferably R^{12} and R^{13} are independently hydrogen or methyl.

Most preferably R^{12} and R^{13} are hydrogen.

Preferably Ar^1 is of the formula (A) or (B).

- 10 Preferably R^6 is hydrogen, C_{1-4} alkyl, hydroxy C_{1-4} alkyl, amino C_{1-4} alkyl, fluoro C_{1-4} alkyl, difluoro C_{1-4} alkyl, C_{1-4} alkoxy or C_{1-4} alkoxy C_{1-4} alkyl.

More preferably R^6 is hydrogen, methyl, fluoromethyl, difluoromethyl, methoxy or methoxymethyl.

Most preferably R^6 is hydrogen or methyl.

- 15 Preferably m is 0 or 1.

Preferably R^5 is hydrogen or methyl.

More preferably R^5 is hydrogen.

In a particular aspect Ar^1 is 1-methylimidazol-5-yl.

- 20 Preferred heteroaryl value for Ar^2 are thiazolyl, pyridyl, triazolyl, pyrimidyl, pyrazinyl or pyridazinyl, especially thiazol-2-yl.

When Ar² is phenyl, it is preferably unsubstituted or monosubstituted.

In one aspect, when Ar² is phenyl, it is unsubstituted.

In another aspect when Ar² is phenyl, it is monosubstituted in the para position.

Preferred substituents for ring carbon atoms in Ar² include C₁₋₄alkyl, halo, nitro,
5 cyano and C₁₋₄alkoxyC₁₋₄alkyl.

More preferred substituents for ring carbon atoms in Ar² include methyl, ethyl, fluoro, chloro, cyano, methoxymethyl and ethoxyethyl.

When Ar² is phenyl it is preferably substituted by fluoro.

When Ar² is thiazolyl it is preferably unsubstituted.

10 Preferably Ar² is 4-fluorophenyl or thiazolyl.

Most preferably Ar² is 4-fluorophenyl or thiazol-2-yl.

Preferably Ar³ is phenyl or pyridyl.

Most preferably Ar³ is phenyl.

15 Preferably, when n is 0, Ar³ is substituted by R² in the 4-position and -(CH₂)_nR³ in the 3- or 5-position and when n is 1 or 2, Ar³ is substituted by R² in the 3- or 5-position and -(CH₂)_nR³ in the 4-position. The positions indicated are relative to the point of attachment of Ar³ to -(CH₂)_p.


Preferably n is 0 or 2.

In a particular aspect n is 0.

20 In one aspect p is 0.

In another aspect p is 1.

R² is preferably a group of formula:



R⁷ is preferably hydrogen or methyl, especially hydrogen. In R⁸, q is preferably 1-4, more
25 preferably 1 or 2, especially 2.

Within R⁸, R¹⁰ is preferably C₁₋₄alkylsulfanyl, C₁₋₄alkylsulfinyl, C₁₋₄alkylsulfonyl, hydroxy or C₁₋₄alkoxy. More preferably R¹⁰ is methylsulfanyl or methylsulfonyl.

R⁹ is preferably hydroxy, C₁₋₄alkoxy, C₃₋₉cycloalkyloxy, heterocyclyloxy or heterocyclylC₁₋₄alkoxy. More preferably R⁹ is hydroxy, methoxy, propoxy, butoxy, tert-

butoxy, cyclopentyloxy, piperidin-4-yloxy or morpholinoC₁₋₄alkyl. Most preferably, R⁹ is methoxy, propoxy, butoxy, tert-butoxy or cyclopentyloxy.

Preferably R¹¹ in R⁹ is phenyl.

Preferred substituents for NH groups in heterocyclic groups in R⁹ include methyl, ethyl, acetyl, propionyl, fluoromethyl, difluoromethyl and trifluoromethyl.

More preferred substituents for NH groups in heterocyclic groups in R⁹ include methyl and acetyl.

Preferred substituents for ring carbon atoms in phenyl or heteroaryl groups in R¹¹ include methyl, halo, C₁₋₄alkanoyl, nitro, cyano, C₁₋₄alkylsulfinyl, C₁₋₄alkylsulfonyl, carbamoyl, C₁₋₄alkylcarbamoyl and diC₁₋₄alkylcarbamoyl.

Preferably R³ is phenyl, pyridyl or thiazolyl.

Most preferably R³ is phenyl.

Preferred substituents for ring carbon atoms in R³ include C₁₋₄alkyl, halo, C₁₋₄alkoxy, nitro, cyano and C₁₋₄alkoxyC₁₋₄alkyl.

More preferred substituents for ring carbon atoms in R³ include methyl, fluoro, chloro, methoxy, nitro, cyano and methoxymethyl.

A preferred substituent for a ring NH group in a heteroaryl group in R³ is C₁₋₄alkyl, particularly methyl.

When R³ is phenyl it is preferably substituted in the 4-position.

Preferably n is 0 or 2.

A preferred compound of the invention is a compound of the Formula (I) wherein :

Ar¹ is of the formula (A) or (B);

R⁵ is hydrogen or methyl;

R⁶ is hydrogen, C₁₋₄alkyl, fluoroC₁₋₄alkyl, difluoroC₁₋₄alkyl, C₁₋₄alkoxy or

C₁₋₄alkoxyC₁₋₄alkyl;

m is 0 or 1;

R¹² and R¹³ are independently hydrogen or methyl;

Ar² is phenyl or thiazolyl;

Ar³ is phenyl or pyridyl, the ring being substituted on ring carbon atoms by R² and -(CH₂)_nR³ and wherein Ar³ is attached to Ar¹C(R¹²)R¹³CH(Ar²)O- by a ring carbon atom; and

- 9 -

n is 0, 1 or 2;

R² is of the formula (2) wherein R⁷ is hydrogen or methyl;

R⁸ is $-(CH_2)_qR^{10}$ wherein q is 0-4 and R¹⁰ is C₁₋₄alkylsulfanyl, C₁₋₄alkylsulfinyl, C₁₋₄alkylsulfonyl, hydroxy or C₁₋₄alkoxy;

5 R⁹ is hydroxy, C₁₋₄alkoxy, C₃₋₉cycloalkyloxy, heterocycloxy or heterocyclylC₁₋₄alkoxy;

or R² is of the formula (3);

R³ is phenyl, pyridyl or thiazolyl; and phenyl, heteroaryl and heterocyclyl rings in R³, R⁹ and Ar² are independently optionally substituted on ring carbon atoms by one or two
 10 substituents selected from C₁₋₄alkyl, halo, C₁₋₄alkoxy, C₁₋₄alkanoyl, nitro, cyano, C₁₋₄alkylsulfinyl, C₁₋₄alkylsulfonyl, carbamoyl, C₁₋₄alkylcarbamoyl and diC₁₋₄alkylcarbamoyl; and optionally substituted on ring NH groups by C₁₋₄alkyl, C₁₋₄alkanoyl, fluoromethyl, difluoromethyl or trifluoromethyl;
 or a pharmaceutically-acceptable salt, prodrug or solvate thereof.

15 A more preferred compound of the invention is a compound of the formula (I) wherein:

Ar¹ is of the formula (A) or (B);

R⁵ is hydrogen or methyl;

R⁶ is hydrogen, methyl, fluoromethyl, difluoromethyl, methoxy or methoxymethyl;

20 m is 0 or 1;

R¹² and R¹³ are independently hydrogen or methyl;

Ar² is phenyl or thiazolyl, optionally substituted on ring carbon atoms by one or two substituents selected from C₁₋₄alkyl, halo, nitro, cyano and C₁₋₄-alkoxyC₁₋₄alkyl;

Ar³ is phenyl or pyridyl; the ring being substituted on ring carbon atoms by R² and
 25 $-(CH_2)_nR^3$ and wherein Ar³ is attached to Ar¹C(R¹²)R¹³CH(Ar²)O- by a ring carbon atom; and n is 0, 1 or 2;

R² is of formula (2) wherein R⁷ is hydrogen or methyl;

R⁸ is $-(CH_2)_qR^{10}$ wherein q is 1 or 2, and

R¹⁰ is methylsulfanyl or methylsulfonyl;

30 R⁹ is hydroxy, methoxy, propoxy, butoxy, tert-butoxy, cyclopentyloxy, piperidin-4-yloxy, or morpholinoC₁₋₄alkyl; or R² is of the formula (3);

R³ is phenyl optionally substituted by one or two substituents selected from C₁₋₄alkyl, halo, C₁₋₄alkoxy, nitro, cyano and C₁₋₄alkoxyC₁₋₄alkyl; or a pharmaceutically-acceptable salt, prodrug or solvate thereof.

An even more preferred compound of the invention is a compound of the formula (I)

5 wherein:

Ar¹ is of the formula (A) or (B);

R⁵ is hydrogen or methyl;

R⁶ is hydrogen or methyl;

m is 0 or 1;

10 R¹¹ and R¹² are hydrogen;

Ar² is phenyl or thiazol-2-yl wherein the phenyl ring is optionally substituted by fluoro;

Ar³ is phenyl; the ring being substituted on ring carbon atoms by R² and -(CH₂)_nR³ and wherein Ar³ is attached to Ar¹(R¹²)R¹³CH(Ar²)O- by a ring carbon atom;

15 and n is 0, 1 or 2;

R² is of the formula (2) wherein R⁷ is hydrogen;

R⁸ is -(CH₂)_qR¹⁰ wherein q is 2 and

R¹⁰ is methylsulfanyl or methylsulfonyl;

R⁹ is hydroxy, methoxy, propoxy, butoxy, tert-butoxy, cyclopentyloxy, piperidin-4-
20 yloxy, or 2-morpholinoprop-2-yl;

R³ is phenyl optionally substituted by fluoro;

or a pharmaceutically-acceptable salt, prodrug or solvate thereof.

Particular compounds of the present invention include:


methyl (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-
25 fluorophenyl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
(2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-
fluorophenyl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid;
tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-
fluorophenyl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
30 methyl (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(thiazol-2-
yl)ethoxymethyl]benzoylamino}-4-methylsulfanyl butyric acid;

- (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(thiazol-2-yl)ethoxymethyl]benzoylamino}-4-methylsulfanyl butyric acid;
 methyl (2S)-2-{2-(4-fluorophenethyl)-5-[2-(imidazol-1-yl)-1-(4-fluorophenyl)-ethoxymethyl]benzoylamino}-4-methylsulfanyl butyrate;
- 5 (2S)-2-{2-(4-fluorophenethyl)-5-{2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid;
 methyl (2S)-2-{2-(4-(fluorobenzyl)-5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
 (2S)-2-{2-(4-fluorobenzyl)-5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid;
- 10 methyl (2S)-2-{2-phenyl-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate; or
 (2S)-2-{2-phenyl-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid;
- 15 methyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-2-methyl-4-methylsulfanylbutyrate;
 (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-2-methyl-4-methylsulfanylbutyric acid;
 N-(4-chlorobenzenesulfonyl)-(2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyramide;
- 20 2-(morpholinomethyl)prop-2-yl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
 methyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate;
- 25 (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyric acid;
 tert-butyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate;
 cyclopentyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate;
- 30

- tert-butyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid;
- 5 methyl (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
(2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid;
- tert-butyl (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
- 10 tert-butyl (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
(2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid;
- 15 methyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
(2S)-2-{5-[1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid;
- tert-butyl (2S)-2-{4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
- 20 tert-butyl (2S)-2-{4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
2-{4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid;
- 25 methyl (2S)-2-{5-[2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
tert-butyl (2S)-2-{5-[2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
(2S)-2-{5-[2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid;
- 30


- tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyrate;
- tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfonylbutyrate;
- 5 tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
- tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate;
- (2S)-2-{2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]pyrid-3-
10 oylamino}-4-methylsulfanylbutyric acid;
- tert-butyl (2S)-2-{2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]pyrid-3-oilamino}-4-methylsulfanylbutyrate;
- tert-butyl (2S)-2-{3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyrid-2-oilamino}-4-methylsulfanylbutyrate;
- 15 cyclopentyl (2S)-2-{3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyrid-2-oilamino}-4-methylsulfanylbutyrate;
- (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyric acid;
- (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-
20 yl)ethoxy]benzoylamino}-4-methylsulfonylbutyric acid;
- tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
- tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate;
- 25 tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate; and
- tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate;
- and pharmaceutically-acceptable salts thereof.
- 30 In another aspect the invention provides an inhibitor of ras farnesylation of Formula (1):

- 14 -



Formula (1)

wherein Ar¹ represents:




Ar² is phenyl or heteroaryl;

p is 0 or 1;

Ar³ is phenyl, pyridinyl, pyridazinyl, pyrimidyl or pyrazinyl, the ring being substituted on 5 ring carbon atoms by R² and $-(\text{CH}_2)_n\text{R}^3$ and wherein Ar³ is attached to Ar¹CH₂CH(Ar²)O- by a ring carbon atom;

R² is a group of the Formula (2):




Formula (2)

wherein R⁷ is hydrogen or C₁₋₄alkyl, R⁸ is $-(\text{CH}_2)_q\text{R}^{10}$ wherein q is 0-4 and R¹⁰ is C₁₋

10 ₄alkylsulfanyl, C₁₋₄alkylsulfinyl, C₁₋₄alkylsulfonyl, hydroxy, C₁₋₄alkoxy, carbamoyl, N-C₁₋₄alkyl carbamoyl, N,N-(diC₁₋₄alkyl)carbamoyl, C₁₋₄alkyl, phenyl, thienyl, or C₁₋₄alkanoylamino, R⁹ is hydroxy, C₁₋₄alkoxy, C₃₋₉cycloalkyloxy, heterocyclyloxy, heterocyclylC₁₋₄alkoxy or -NH-SO₂-R¹¹ wherein R¹¹ represents, trifluoromethyl, C₁₋₄alkyl, phenyl, heteroaryl, arylC₁₋₄alkyl or heteroarylC₁₋₄alkyl;

15 or R² represents a lactone of Formula (3)

- 15 -



Formula (3)

the group of Formula (2) or (3) having L or D configuration at the chiral alpha carbon in the corresponding free amino acid;

n is 0, 1 or 2;

5 R^3 is phenyl or heteroaryl;

R^3 and Ar^2 are independently optionally substituted by up to three substituents selected from C_{1-4} alkyl, halogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, amino, C_{1-4} alkylamino, di(C_{1-4} alkyl)amino, C_{1-4} alkanoylamino, nitro, cyano, carboxy, C_{1-4} alkoxycarbonyl, thiol, C_{1-4} alkylsulfanyl, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkanesulphonamido, N-(C_{1-4} alkylsulphonyl)-N- C_{1-4} alkylamino, aminosulfonyl, N-(C_{1-4} alkyl)aminosulfonyl, N,N-di(C_{1-4})aminosulfonyl, carbamoyl, N-(C_{1-4} alkyl)carbamoyl, N,N-(di C_{1-4} alkyl)carbamoyl, carbamoyl C_{1-4} alkyl, N-(C_{1-4} alkyl)carbamoyl C_{1-4} alkyl, N,N-(di C_{1-4} alkyl)carbamoyl C_{1-4} alkyl, hydroxy C_{1-4} alkyl and C_{1-4} alkoxy C_{1-4} alkyl;

10 R^5 is hydrogen, C_{1-4} alkyl, aryl C_{1-4} alkyl;

15 R^6 is hydrogen, C_{1-4} alkyl, hydroxy C_{1-4} alkyl, sulfanyl C_{1-4} alkyl, N-(C_{1-4} alkyl)amino C_{1-4} alkyl or aryl C_{1-4} alkyl; m is 0, 1 or 2;

or a pharmaceutically-acceptable salt, prodrug or solvate thereof.

In one aspect p is 0.

In another aspect p is 1.

20 Compounds of Formula (1) may form salts which are within the ambit of the invention. Pharmaceutically acceptable salts are preferred although other salts may be useful in, for example, isolating or purifying compounds.

When the compound contains a basic moiety it may form pharmaceutically-acceptable salts with a variety of inorganic or organic acids, for example hydrochloric, hydrobromic,

25 sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. A suitable pharmaceutically-acceptable salt of the invention when the compound contains an acidic moiety is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic

base which affords a pharmaceutically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

Solvates, for example hydrates, are also within the ambit of the invention and may be
5 prepared by generally known methods.

Various forms of prodrugs are well known in the art. For examples of such prodrug derivatives, see:

- a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);
- 10 b) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen and H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard p. 113-191 (1991);
- c) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);
- d) H. Bundgaard, *et al.*, Journal of Pharmaceutical Sciences, 77, 285 (1988); and
- 15 e) N. Kakeya, *et al.*, Chem Pharm Bull, 32, 692 (1984).

Examples of pro-drugs include *in vivo* hydrolysable esters of a compound of the Formula I. Suitable pharmaceutically-acceptable esters for carboxy include C₁₋₈alkyl esters, C₅₋₈cycloalkyl esters, cyclic amine esters, C₁₋₆alkoxymethyl esters for example methoxymethyl, C₁₋₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl
20 esters, C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl wherein alkyl, cycloalkyl and cyclicamino groups are optionally substituted by, for example, phenyl, heterocyclcyl, alkyl, amino, alkylamino, dialkylamino,
25 hydroxy, alkoxy, aryloxy or benzyloxy, and may be formed at any carboxy group in the compounds of this invention.

According to another aspect of the invention there is provided a pharmaceutical composition comprising a compound as defined in Formula (1) or an individual compound listed above together with a pharmaceutically-acceptable diluent or carrier. A preferred
30 pharmaceutical composition is in the form of a tablet.

The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for
5 example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using
10 conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

Suitable pharmaceutically-acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium
15 carbonate, granulating and disintegrating agents such as corn starch or algenic acid; binding agents such as starch; lubricating agents such as magnesium stearate, stearic acid or talc; preservative agents such as ethyl or propyl p-hydroxybenzoate, and anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the
20 gastrointestinal tract, or to improve their stability and/or appearance, in either case, using conventional coating agents and procedures well known in the art.

Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with
25 water or an oil such as peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions generally contain the active ingredient in finely powdered form together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation
30 products of an alkylene oxide with fatty acids (for example polyoxethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols, for example

heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters
5 derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl *p*-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening
10 agents (such as sucrose, saccharine or aspartame).

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring
15 agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or
20 wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients such as sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil,
25 or a mineral oil, such as for example liquid paraffin or a mixture of any of these. Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, an esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene
30 sorbitan monooleate. The emulsions may also contain sweetening, flavouring and preservative agents.

Syrups and elixirs may be formulated with sweetening agents such as glycerol, propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

The pharmaceutical compositions may also be in the form of a sterile injectable
5 aqueous or oily suspension, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which have been mentioned above. A sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol.

10 Suppository formulations may be prepared by mixing the active ingredient with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Suitable excipients include, for example, cocoa butter and polyethylene glycols.

Topical formulations, such as creams, ointments, gels and aqueous or oily solutions or
15 suspensions, may generally be obtained by formulating an active ingredient with a conventional, topically acceptable, vehicle or diluent using conventional procedure well known in the art.

Compositions for administration by insufflation may be in the form of a finely divided powder containing particles of average diameter of, for example, 30 μ or much less, the
20 powder itself comprising either active ingredient alone or diluted with one or more physiologically acceptable carriers such as lactose. The powder for insufflation is then conveniently retained in a capsule containing, for example, 1 to 50mg of active ingredient for use with a turbo-inhaler device, such as is used for insufflation of the known agent sodium cromoglycate.

25 Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

For further information on Formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient. For further information on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The size of the dose for therapeutic or prophylactic purposes of a compound of the Formula (1) will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well known principles of medicine. As mentioned above, compounds of the Formula (1) are useful in treating diseases or medical conditions which are due alone or in part to the effects of farnesylation of ras.

In using a compound of the Formula (1) for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for example, 0.5 mg to 75 mg per kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous administration, a dose in the range, for example, 0.5 mg to 30 mg per kg body weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.5 mg to 25 mg per kg body weight will be used. Oral administration is however preferred.

Compounds of this invention may be useful in combination with known anti-cancer and cytotoxic agents. If formulated as a fixed dose such combination products employ the compounds of this invention within the dosage range described herein and the other

pharmaceutically active agent within its approved dosage range. Sequential use is contemplated when a combination formulation is inappropriate.

According to another aspect of the invention there is provided a compound of Formula (1) or a pharmaceutically-acceptable salt thereof, for use as a medicament.

5 According to another aspect of the invention there is provided a compound of Formula (1) or a pharmaceutically-acceptable salt thereof, for use in preparation of a medicament for treatment of a disease mediated through farnesylation of ras.

According to another aspect of the present invention there is provided a method of treating ras mediated diseases, especially cancer, by administering an effective amount of a
10 compound of Formula (1) or a pharmaceutically-acceptable salt thereof, to a mammal in need of such treatment.

Diseases or medical conditions may be mediated alone or in part by farnesylated ras. A particular disease of interest is cancer. Specific cancers of interest include:

- carcinoma, including that of the bladder, breast, colon, kidney, liver, lung,
15 ovary, pancreas, stomach, cervix, thyroid and skin;
- hematopoietic tumors of lymphoid lineage, including acute lymphocytic leukemia, B-cell lymphoma and Burkett's lymphoma;
- hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous leukemias and promyelocytic leukemia;
- 20 - tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; and
- other tumors, including melanoma, seminoma, teratocarcinoma, neuroblastoma and glioma.


The compounds of Formula (1) are especially useful in treatment of tumors having a
25 high incidence of ras mutation, such as colon, lung, and pancreatic tumors. By the administration of a composition having one (or a combination) of the compounds of this invention, development of tumors in a mammalian host is reduced.

Compounds of Formula (1) may also be useful in the treatment of diseases other than cancer that may be associated with signal transduction pathways operating through Ras, e.g.,
30 neuro-fibromatosis.

Compounds of Formula (1) may also be useful in the treatment of diseases associated with CAAX-containing proteins other than Ras (e.g., nuclear lamins and transducin) that are also post-translationally modified by the enzyme farnesyl protein transferase.

Although the compounds of the Formula (1) are primarily of value as therapeutic agents for use in warm-blooded animals (including man), they are also useful whenever it is required to inhibit the effects of activation of ras by farnesylation. Thus, they are useful as pharmacological standards for use in the development of new biological tests and in the search for new pharmacological agents.

In another aspect the present invention provides a process for preparing a compound of the Formula (1) or a pharmaceutically-acceptable salt prodrug or solvate thereof which process comprises:
deprotecting a compound of the formula (4)



(4)

wherein Ar^{1'} is Ar¹ or protected Ar¹, Ar^{2'} is Ar² or protected Ar² and Ar^{3'} is Ar³ or protected Ar³; wherein at least one protecting group is present; and thereafter if necessary:

- (i) forming a pharmaceutically-acceptable salt,
- (ii) forming a prodrug,
- (iii) forming a solvate.

Protecting groups may in general be chosen from any of the groups described in the literature or known to the skilled chemist as appropriate for the protection of the group in question, and may be introduced by conventional methods.

Protecting groups may be removed by any convenient method as described in the literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so as to effect removal of the protecting group with minimum disturbance of groups elsewhere in the molecule.

Specific examples of protecting groups are given below for the sake of convenience, in which "lower" signifies that the group to which it is applied preferably has 1-4 carbon atoms.

It will be understood that these examples are not exhaustive. Where specific examples of methods for the removal of protecting groups are given

below these are similarly not exhaustive. The use of protecting groups and methods of deprotection not specifically mentioned is of course within the scope of the invention.

5 A carboxy protecting group may be the residue of an ester-forming aliphatic or araliphatic alcohol or of an ester-forming silanol (the said alcohol or silanol preferably containing 1-20 carbon atoms).

Examples of carboxy protecting groups include straight or branched chain C₁₋₁₂alkyl groups (for example isopropyl, *t*-butyl); lower alkoxy lower alkyl groups (for example
10 methoxymethyl, ethoxymethyl, isobutoxymethyl); lower aliphatic acyloxy lower alkyl groups, (for example acetoxymethyl, propionyloxymethyl, butyryloxymethyl, pivaloyloxymethyl); lower alkoxycarbonyloxy lower alkyl groups (for example 1-methoxycarbonyloxyethyl, 1-ethoxycarbonyloxyethyl); phenyl lower alkyl groups (for example benzyl, *p*-methoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, benzhydryl and phthalidyl); tri(lower
15 alkyl)silyl groups (for example trimethylsilyl and *t*-butyldimethylsilyl); tri(lower alkyl)silyl lower alkyl groups (for example trimethylsilylethyl); and C₂₋₆alkenyl groups (for example allyl and vinyloethyl).

Methods particularly appropriate for the removal of carboxy protecting groups include for example acid-, base-, metal- or enzymically-catalysed hydrolysis.

20 Examples of hydroxy protecting groups include lower alkyl groups (for example *t*-butyl), lower alkenyl groups (for example allyl); lower alkanoyl groups (for example acetyl); lower alkoxycarbonyl groups (for example *t*-butoxycarbonyl); lower alkenyloxycarbonyl groups (for example allyloxycarbonyl); phenyl lower alkoxycarbonyl groups (for example benzoyloxycarbonyl, *p*-methoxybenzyloxycarbonyl,
25 *o*-nitrobenzyloxycarbonyl, *p*-nitrobenzyloxycarbonyl); tri lower alkylsilyl (for example trimethylsilyl, *t*-butyldimethylsilyl) and phenyl lower alkyl (for example benzyl) groups.

Examples of amino protecting groups include formyl, aralkyl groups (for example benzyl and substituted benzyl, *p*-methoxybenzyl, nitrobenzyl and 2,4-dimethoxybenzyl, and
30 triphenylmethyl); di-*p*-anisylmethyl and furylmethyl groups; lower alkoxycarbonyl (for example *t*-butoxycarbonyl); lower alkenyloxycarbonyl (for example allyloxycarbonyl); phenyl

lower alkoxy carbonyl groups (for example benzyloxycarbonyl, p-methoxybenzyloxycarbonyl, o-nitrobenzyloxycarbonyl, p-nitrobenzyloxycarbonyl; trialkylsilyl (for example trimethylsilyl and t-butyldimethylsilyl); alkylidene (for example methylenedene); benzylidene and substituted benzylidene groups.



- 5 Methods appropriate for removal of hydroxy and amino protecting groups include, for example, acid-, base-, metal- or enzymically-catalysed hydrolysis, for groups such as p-nitrobenzyloxycarbonyl, hydrogenation and for groups such as o-nitrobenzyloxycarbonyl, photolytically.

The reader is referred to Advanced Organic Chemistry, 4th Edition, by Jerry March,
10 published by John Wiley & Sons 1992, for general guidance on reaction conditions and reagents. The reader is referred to Protective Groups in Organic Synthesis, 2nd Edition, by Green *et al.*, published by John Wiley & Sons for general guidance on protecting groups.

Compounds of the formula (1) and (4) can be formed by :


- (i) reacting a compound of the formula (5) with a compound of the formula (6)

15



or (iii) converting one value of R⁹ in R² into another value of R⁹;

- or (iii) reacting a compound in which R² in Ar³ is carboxy with a compound of the formula
20 (7):



wherein p, Ar¹, Ar², Ar³, R⁷ and R⁸ are as hereinabove defined, R²¹ is R⁹ or a carboxy protecting group and when p is 1, L is a leaving group, and when p is 0, L is hydroxy; and thereafter if necessary:


- 25 (i) removing any protecting groups;

(ii) forming a pharmaceutically-acceptable salt, prodrug or solvate thereof.

When p is 1, compounds of the formula (5) and (6) are conveniently reacted together in the presence of a base such as sodium hydride, butyl lithium or potassium tert-butoxide, in an aprotic solvent such as tetrahydrofuran (THF), dimethyl formamide (DMF) or
 5 dimethylacetamide (DMA), at a non-extreme temperature for example 0 °C to ambient temperature. L is preferably halo, mesyloxy or tosyloxy.


When p is 0, a compound of the formula (5) and a compound of the formula (6) are conveniently reacted together under conditions known for the Mitsunobu reaction. This typically involves reacting the reagents together in the presence of di(C₁-
 10 4alkyl)azocarboxylate or 1', 1'-(azodicarbonyl)dipiperidine and a phosphorus reagent such as tributylphosphine or triphenylphosphine in an inert solvent such as toluene, benzene, tetrahydrofuran (THF) or diethylether, at non-extreme temperatures such as in the range -20°C to ambient temperature (see Progress in the Mitsunobu Reaction. A Review, David L. Hughes, Organic Preparations and Procedures Int., 28 (2), 127-164 (1996)).

15 A compound of the formula (5) can be prepared by reducing a compound of the formula (8):




wherein Ar^{1'} and Ar^{2'} are as hereinabove defined. Suitable reducing agents include sodium borohydride or lithium aluminium hydride. Typically, when sodium borohydride is the
 20 reducing agent, an alcohol is used as solvent in a temperature range of ambient temperature to 60 °C, and when lithium hydride is used diethyl ether or THF are used as solvents.

A compound of the formula (8) can be prepared by introducing Ar^{1'} into a compound of the formula (9):



wherein Ar^{2'} is as hereinabove defined and L¹ is a leaving group such as mesyloxy, tosyloxy, triflate or halo, preferably bromo. The reaction is conveniently carried out in the presence of a base such as sodium hydride, sodium hydroxide, butyl lithium or potassium carbonate. In some cases a base may not be necessary.

- 5 A compound of the formula (9) is conveniently formed from a compound of the formula (10):




wherein Ar^{2'} is as hereinabove defined.

The compound of the formula (10) may be converted to a compound in which L¹ is
 10 bromo by bromination with, for example, N-bromosuccinimide, carbon tetrabromide or bromine or to a compound in which L¹ is chloro by chlorination with for example chlorine. When L¹ is mesyloxy or tosyloxy by oxidising the compound of the formula (10) to an alcohol and converting the hydroxy group to mesyloxy or tosyloxy using a meyl halide or tosyl halide.

- 15 When p is 1, a compound of the formula (6) is typically formed by introducing a leaving group into a compound of the formula Ar^{3'}-CH₃. When L is bromo Ar^{3'}-CH₃ can be brominated using for example N-bromosuccinimide, carbon tetrabromide or bromine. When L is chloro, a chlorinating agent such as chlorine could be used and when L is mesyloxy or tosyloxy, the methyl group in Ar^{3'}-CH₃ is generally oxidised to the alcohol (or oxidised to the
 20 carboxylic acid and then reduced to the alcohol) and the hydroxy group converted to mesyloxy or tosyloxy with, for example, mesyl chloride or tosyl chloride.

The compound of the formula Ar^{3'}-CH₃ could be formed by introducing -(CH₂)_nR³ into a compound of the formula (11):

- 27 -



wherein $R^{2'}$ is as hereinabove defined, A is phenyl, pyridyl, pyrazinyl, pyrimidinyl or pyridazinyl and L^2 is a leaving group.

- 5 When n is 0 and R^3 is phenyl, the compound of the formula (11) is conveniently reacted with phenyl boronic acid in the presence of a palladium catalyst such as palladium tetrakis (triphenylphosphine) palladium (0) under conditions known for the Suzuki reaction (Synth. Commun. 11, 513 (1981)). An aprotic organic solvent such as dimethyl ether (DME), dimethylsulphoxide (DMSO) or THF is generally used and a base such as sodium
- 10 bicarbonate, sodium carbonate and sometimes sodium hydroxide. A fluoride such as caesium fluoride could be used instead of the base (J. Org. Chem. 1994, 59, 6095-6097). Preferably L^2 is bromo or triflate.

- When n is 1 and R^3 is phenyl, the compound of the formula (11), bromo or chloro, is conveniently reacted with a benzylzinc chloride or a benzyl-magnesium bromide in the
- 15 presence of a nickel or palladium catalyst, such as bis(triphenylphosphine)palladium (II) chloride or $Pd_2(dba)_3$, in an inert organic solvent such as tetrahydrofuran (THF). For example see the conditions used for the 'Nagishi' reaction (J. Org. Chem. 42 (10), 1821-1822, 1977).

- When n is 2 and R^3 is phenyl, the compound of the formula (11) is conveniently
- 20 reacted with a styrene under conditions known for the Heck reaction. Briefly this involves an inorganic or organic base such as triethylamine, a palladium catalyst such as bis (o-tolylphosphine)palladium (II) chloride in water. (Acc. Chem. Res. 12, 146-151 (1979), J. Organometallic Chem. 486, 259-262 (1995)).


- The resulting alkene can then be reduced using standard methods known in the art, for
- 25 example, catalytic hydrogenation.

Alternatively the alkyne could be formed by reacting a compound of the formula (11) wherein L^2 is triflate or bromo with a phenyl acetylene in the presence of an organic base such

as triethylamine and a palladium catalyst such as palladium tetrakis (triphenylphosphine). For example see the conditions used for the Sonogashira reaction (J. Org. Chem. 1993, 58, 6614-6619).

The resultant alkyne can be reduced using standard methods known in the art, for example, catalytic hydrogenation.

When p is 0, the compound of the formula (6) can be formed by introducing $-(CH_2)_nR^3$ into a compound of the formula (12).



10 wherein R^2 , A and L^2 are as hereinbefore defined, and P^1 is a hydroxy-protecting group.

When n is 0 and R^3 is phenyl, the compound of the formula (12) is conveniently reacted with phenyl boronic acid in the presence of a palladium catalyst as described above for when p is 1.

When p is 0, L^2 is preferably bromo.

15 When n is 1, and R^3 is phenyl, the compound of the formula (12) wherein L^2 is preferably bromo or chloro, is conveniently reacted with benzylzinc chloride or benzylmagnesium bromide under similar conditions to those described above for when p is 1. When n is 2, the compound of the formula (12) is conveniently reacted with styrene under conditions known for the Heck reaction.

20 The resulting alkene can then be reduced using standard methods known in the art, for example, catalytic hydrogenation.


Alternatively, the alkyne could be formed by reacting a compound of the formula (12) wherein L^2 is triflate or bromo with a phenyl acetylene in the presence of an organic base such as triethylamine and a palladium catalyst such as palladium tetrakis (triphenylphosphine). For example see the conditions used for the Sonogashira reaction (J. Org. Chem. 1993, 58, 6614-6619).

The resultant alkyne can be reduced using standard methods known in the art, for example, catalytic hydrogenation.


The protecting group P^1 can then be removed to leave a compound of the formula (6).

- 29 -

A compound of the formula (1) or (4) could be prepared via a sequence of steps from a compound of the formula (12).



- 5 When p is 1, a compound of the formula (1) or (4) could be prepared via a sequence of steps from a compound of the formula (13):



wherein L² and A are as hereinabove defined and P is a carboxy protecting group.

L² can be replaced with the group -(CH₂)_nR³ using the methodology described above.

- 10 The methyl group could then be converted to a -CH₂L group and the resultant compound reacted with a compound of the formula (5). The carboxy group in the resultant product can then be deprotected and reacted with the appropriate amino acid derivative to form R², under conditions described hereinbelow, and hence a compound of the formula (1) or (4).

- When p is 0, a compound of the formula (1) or (4) could be prepared from a
15 compound similar to that of formula (13) but wherein the methyl group is replaced by a protected-hydroxy group. L² can then be converted to -(CH₂)_nR³, the hydroxy group removed and the resultant compound reacted with a compound of the formula (5). Subsequent steps are also similar to those described above for when p is 1.

- A compound of the formula (1) in which R⁹ in R² is alkoxy can conveniently be
20 hydrolysed to another compound of the formula (1) in which R⁹ is hydroxy using standard methods known in the art. For example, the alkoxy group could be subjected to acid or base hydrolysis with, for example, in the case of base hydrolysis, aqueous sodium hydroxide solution in an organic solvent such as an alcohol in a temperature range of ambient temperature to 60°C. When R⁹ is a hydroxy group the carboxy group in a compound of the

formula (1) can be converted to an acylsulphonamide by reacting the carboxy group with the appropriate sulphonamido group in the presence of an organic base such as triethylamine or dimethylaminopyridine, in an inert organic solvent such as dimethylformamide (DMF), in temperature range of -20°C to ambient temperature.

5 The reaction between a compound in which R² in Ar^{3'} is carboxy and a compound of the formula (7) is generally carried out in the presence of a reagent that converts the carboxy group into a reactive ester, for example a carbodiimide such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC) or pentafluorophenyl, and in the presence of an organic base such as N-methylmorpholine or dimethylaminopyridine (DMAP). The reaction is usually carried out
10 in the temperature range of -20°C to ambient temperature. The reagent, 1-hydroxybenzotriazole, is often added to assist the reaction (see Chem. Ber. 103, 788, 2024 (1970), J. Am. Chem. Soc. 93, 6318 (1971), Helv. Chim. Acta. 56, 717, (1973)). Suitable solvents include DMF and dichloromethane.

A compound of the formula (1) in which R² in Ar^{3'} is carboxy can be prepared by
15 reacting a compound of the formula (5) with a compound of the formula (6) wherein R² in Ar^{3'} is protected carboxy and subsequently removing the protecting group.

Optionally substituents in a compound of the formula (1) and (4) or intermediates in their preparation may be converted into other optional substituents. For example an alkylthio group may be oxidised to an alkylsulphinyl or alkylsulphonyl group, a nitro group reduced to
20 an amino group, a hydroxy group alkylated to a methoxy group, or a bromo group converted to an alkylthio group.

Various substituents may be introduced into compounds of the formulae (1) and (4) and intermediates in this preparation, when appropriate, using standard methods known in the art. For example, an acyl group or alkyl group may be introduced into an activated benzene
25 ring using Friedel-Crafts reactions, a formyl group by formylation with titanium tetrachloride and dichloromethyl ethyl ester, a nitro group by nitration with concentrated nitric acid concentrated sulphuric acid and bromination with bromine or tetra(n-butyl)ammonium tribromide.

Alternatively, a compound similar to a compound of the formula (12) but which
30 contains a methyl group instead of -COOP group could be used as the starting material and

both methyl groups oxidised to carboxylic acids, one selectively reduced to an alcohol with a reducing agent such as borane in THF, and the hydroxy converted to a leaving group.

It will be appreciated that, in certain steps in the reaction sequence to compounds of the formula (1), it will be necessary to protect certain functional groups in intermediates in order to prevent side reactions. Deprotection may be carried out at a convenient stage in the reaction sequence once protection is no longer required.

Biological activity was tested as follows:

(i) In-vitro assay

The following stock solutions were used and the assays were conducted in 96 well plates: TRIS Buffer (500mM TRIS, 50mM MgCl₂.6H₂O, pH=8.0); Farnesyl pyrophosphate (6.4mg/ml); Aprotinin (1.9 mg/ml); Ki-ras (0.5mg/ml, stored at -80°C); Acid ethanol (850ml absolute ethanol + 150ml concentrated HCl).

Farnesyl protein transferase (FPT) was partially purified from human placenta by ammonium sulphate fractionation followed by a single Q-SepharoseTM (Pharmacia, Inc) anion exchange chromatography essentially as described by Ray and Lopez-Belmonte (Ray K P and Lopez-Belmonte J (1992) Biochemical Society Transactions 20 494-497). The substrate for FPT was Kras (CVIM C-terminal sequence). The cDNA for oncogenic val12 variant of human c-Ki-ras-2 4B was obtained from the plasmid pSW11-1 (ATCC). This was then subcloned into the polylinker of a suitable expression vector e.g. pIC147. The Kras was obtained after expression in the E. coli strain, BL21. The expression and purification of c-KI-ras-2 4B and the val12 variant in E. coli has also been reported by Lowe et al (Lowe P N et al. J. Biol. Chem. (1991) 266 1672-1678). The farnesyl protein transferase enzyme preparation was stored at -80°C.

The farnesyl transferase solution for the assay contained the following: dithiothreitol (DTT)(0.6ml of 7.7mg/ml), TRIS buffer (0.6ml), aprotinin (0.48ml), distilled water (1.2ml), farnesyl transferase (0.6ml of the crude enzyme preparation prepared as described above), zinc chloride (12µl of 5mM). This was left at ambient temperature for 30 minutes. After this incubation 60µl Ki-ras solution was added and the whole left to incubate for a further 60 minutes prior to use in the assay.

Assays were performed in 96 well plates as follows: 10µl of test compound solution was added to each well. Then 30µl farnesyl transferase solution (above) was added and the

reaction started by addition of 10µl radiolabelled farnesyl pyrophosphate solution. After 20 minutes at 37°C the reaction was stopped with 100µl acid ethanol (as described in Pompliano D L et al (1992) 31 3800-3807). The plate was then kept for 1 hour at 4°C. Precipitated protein was then collected onto glass fibre filter mats (B) using a Tomtec™ cell harvester and 5 tritiated label was measured in a Wallac™1204 Betaplate scintillation counter. Test compounds were added at appropriate concentrations in DMSO (3% final concentration in test and vehicle control).

(ii) Intracellular farnesylation assay

HER313A cells (Grand et al, 1987 Oncogene 3, 305-314) were routinely cultured in 10 Dulbecos Modified Essential Medium (DMEM) plus 10% foetal calf serum (FCS). For the assay HER313A cells were seeded at 200,000 cells/well in a volume of 2.5ml in a 6 well tissue culture plate. After an overnight incubation at 37°C in 10% CO₂ the medium was removed and replaced with methionine-free minimal essential medium (MEM) and the cells incubated as above for 2 hours. After this time the medium was removed and replaced by 15 methionine-free MEM (1ml) and test compound (1-3µl). The plates were then incubated for a further 2 hours as described above and then 30µCi of ³⁵S-methionine added to each well. The plate was then incubated overnight as described above. The medium was then removed and the cells were lysed with lysis buffer (1ml) (composed of 1000ml phosphate buffered saline, 10ml triton X-100, 5g sodium deoxycholate, 1g sodium dodecylsulphate) containing 20 aprotinin (10µl/ml), the plate scrapped and then left for 10 minutes at 4°C. The lysate was then clarified by centrifugation. To 0.8ml of the clarified lysate 80µl of Y13-259 pan-Ras antibody (isolated from the hybridoma - American Tissue Culture Collection Accession Number CRL-1742) (final concentration approximately 1µg/ml, the exact working concentration was optimised for each batch of antibody isolated) and protein G beads (30µl of 25 0.5µg/ml) were added and the mixture incubated overnight with constant agitation. The pellet was then collected by centrifugation, washed and separated by SDS PAGE using a 15% gel. Radioactive bands were detected using a phosphorimager system.

(iii) Morphology and proliferation assay

MIA PaCa 2 cells (American Tissue Culture Collection Accession Number: 30 CRL-1420) were routinely cultured in Dulbecos Modified Essential Medium (DMEM) plus 10% FCS in a 162 cm² tissue culture flask. For the assay the cells were seeded at 16,000

cells/well, in 12 well plates, in DMEM containing 5% charcoal dextran treated stripped FCS (1ml)(obtained from Pierce and Warriner). The cells were then incubated overnight at 37°C in 10% CO₂. Test compound was then added (10µl) and the cells incubated for 6 days as described above. On days 1, 2, 3 and 6 the cells were monitored for signs of morphological change and toxicity. On day 6 the cells were removed from the plate using trypsin/EDTA and counted to determine the proliferation rate.

Although the pharmacological properties of the compounds of the Formula (1) vary with structural change as expected, in general compounds of the Formula (1) possess an IC₅₀ in the above test in the range, for example, 0.0005 to 50µM. Thus by way of example the compound of Example 2 herein has an IC₅₀ of approximately 0.001µM. No physiologically unacceptable toxicity was observed at the effective dose for compounds tested of the present invention.

The invention will now be illustrated in the following non-limiting Examples in which, unless otherwise stated:-

- 15 (i) evaporations were carried out by rotary evaporation in vacuo and work-up procedures were carried out after removal of residual solids by filtration;
- (ii) operations were carried out at ambient temperature, that is in the range 18-25°C and under an atmosphere of an inert gas such as nitrogen or argon;
- (iii) column chromatography (by the flash procedure) and medium pressure liquid chromatography (MPLC) were performed on Merck Kieselgel silica (Art. 9385) or Merck Lichroprep RP-18 (Art. 9303) reversed-phase silica obtained from E. Merck, Darmstadt, Germany or high pressure liquid chromatography (HPLC) C18 reverse phase silica separation;
- (iv) yields are given for illustration only and are not necessarily the maximum attainable;
- 25 (v) the end-products of the Formula (1) have satisfactory microanalyses and their structures were confirmed by nuclear magnetic resonance (NMR) and mass spectral techniques; chemical shift values were measured on the delta scale; the following abbreviations have been used: s, singlet; d, doublet; t or tr, triplet; m, multiplet; br, broad;
- (vi) intermediates were not generally fully characterised and purity was assessed by 30 thin layer chromatographic, HPLC, infra-red (IR) or NMR analysis;

(vii) melting points are uncorrected and were determined using a Mettler SP62 automatic melting point apparatus or an oil-bath apparatus; melting points for the end-products of the Formula (1) were determined after crystallisation from a conventional organic solvent such as ethanol, methanol, acetone, ether or hexane, alone or in admixture;
5 and

(viii) the following abbreviations have been used:-

	ADDP	1,1'-(azodicarbonyl)dipiperidine
	DEAD	diethyl azodicarboxylate
	DMAP	4-dimethylaminopyridine
10	DMF	dimethylformamide
	EDC	1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide
	HOBt	1-hydroxybenzotriazole
	Pd(dppb)Cl ₂	[1,4-bis(diphenylphosphino)butane]palladium(II) dichloride
	MCPBA	m-chloroperoxybenzoic acid
15	NMM	<u>N</u> -methylmorpholine
	TEA	triethylamine
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran

Example 1**Methyl (2S)-2-[2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoylamino]-4-methylsulfanylbutyrate**

A mixture of 2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoic acid (0.522 g ; 1.2 mmol), methyl (2S)-2-amino-4-methylsulfanylbutanoate hydrochloride (L-methionine methyl ester hydrochloride) (0.24 g; 1.2 mmol), HOBT (0.163 g ; 1.2 mmol), EDC (0.23 g ; 1.2 mmol) and N-methylmorpholine (0.132 ml ; 1.2 mmol) in dichloromethane (15 ml) was stirred under an argon atmosphere for 5 hours. After evaporation to dryness, the residue was purified by flash column chromatography eluting with petroleum ether / ethanol (98:2). The resulting compound was dissolved in dichloromethane (2 ml), treated at 0°C with a 3.8N solution of HCl in ether (0.265 ml) and diluted with ether (100 ml) to precipitate the hydrochloride salt which was filtered and dried to give the title compound.

Yield : 75%

¹H NMR (DMSO d₆ + CF₃COOD, 400 MHz) δ : 1.8-1.95 (2H, m) ; 2 (3H, s) ; 2.15-2.35 (2H, m) ; 3.64 (3H, s) ; 4.3-4.55 (5H, m) ; 4.9 (1H, m) ; 7.1-7.7 (13H, m) ; 8.6 (1H, d) ; 9.05 (1H, s).

Anal. Calculated for C₃₁H₃₁F₂N₃O₄S, H₂O, 0.9 HCl

	C 60.44	H 5.28	N 6.82	S 5.20	
20	Found	C 60.07	H 5.39	N 6.69	S 4.95
	MS (ESI) m/z 580 (MH ⁺)				

The starting material was prepared as follows :

Triflic anhydride (170 ml ; 1.01 mol) was added to a solution of methyl 2-hydroxy-4-methylbenzoate (153 g ; 0.92 mol) in pyridine (1.5 l), at 0°C. The mixture was stirred at ambient temperature overnight. After evaporation of the pyridine, the residue was acidified to pH 3.5 with 6N HCl and extracted with ether. The organic phase was evaporated and the residue purified by flash column chromatography eluting with a gradient of 0-5% ethyl acetate / petroleum ether to give methyl 2-trifluoromethylsulfonyloxy-4-methylbenzoate (245 g ; 90%).

¹H NMR (CDCl₃, 400 MHz) δ : 1.55 (3H, s) ; 2.45 (3H, s) ; 3.94 (3H, s) ; 7.09 (1H, s) ; 7.26 (1H, s) ; 7.98 (1H, d).

Tetrakis(triphenylphosphine) palladium (9 g ; 7.8 mmol) and ethanol (780 ml) was added to a suspension of methyl 4-methyl-2-trifluoromethanesulphonyloxybenzoate (58 g ; 0.195 mol), 2M aqueous solution of sodium carbonate (250 ml ; 0.5 mol), 4-fluorophenylboronic acid (30 g ; 0.214 mol) and lithium chloride (16.5 g ; 0.39 mol) in toluene (1.65 ml), under an argon atmosphere. The mixture was refluxed for 4 hours, diluted with ethyl acetate (1 l) and washed with aqueous sodium hydroxide solution 1N (1 l). The organic phase was evaporated and the residue purified by flash column chromatography using ethyl acetate/petroleum ether (95:5) to give methyl 2-(4-fluorophenyl)-4-methylbenzoate (46.8 g ; 99%).

¹H NMR (CDCl₃, 400 MHz) δ : 2.41 (3H, s) ; 3.64 (3H, s) ; 7-7.03 (6H, m) ; 7.7 (1H, d).

A solution of methyl 2-(4-fluorophenyl)-4-methylbenzoate (54.18 g ; 0.22 mol), N-bromosuccinimide (39.6 g ; 0.22 mol), 2,2'-azobis(2-methylproprionitrile) (0.25 g ; 1.5 mmol) and benzoylperoxide (0.25 g ; 1 mmol) in tetrachloromethane (550 ml) was heated at reflux for 6 hours. The solid was filtered and the filtrate evaporated to give methyl 4-bromomethyl-2-(4-fluorophenyl)benzoate as an oil (79.7 g ; 79%) which was used in the next step without purification.

¹H NMR (CDCl₃, 400 MHz) δ : 3.65 (3H, s) ; 4.51 (2H, s) ; 7-7.5 (6H, m) ; 7.82 (1H, m)

To a solution of 1-(4-fluorophenyl)-2-(imidazol-1-yl)ethanol (0.628 g ; 3 mmol) in THF (30 ml) was added sodium hydride, under an argon atmosphere, (0.15 g ; 3.6 mmol). After stirring for 10 minutes, methyl 4-bromomethyl-2-(4-fluorophenyl) benzoate (2 g ; 6.09 mmol) was added. The mixture was stirred at ambient temperature overnight. After evaporation to dryness, the residue was extracted with ethyl acetate and purified by flash column chromatography eluting with dichloromethane/ethanol (97:3) to give methyl 2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-(fluorophenyl)ethoxymethyl)]benzoate as an oil (0.635 g ; 46%).

¹H NMR (CDCl₃, 400 MHz) δ : 3.65 (3H, s) ; 4-4.6 (5H, m) ; 6.87 (1H, s) ; 6.98 (1H, s) ; 7-7.3 (10H, m) ; 7.41 (1H, d) ; 7.80 (1H, d).

Methyl 2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoate (0.635 g ; 1.41 mmol) in methanol (15 ml) was treated with 2N aqueous sodium hydroxide solution (1.77 ml ; 3.54 mmol) at ambient temperature for 8 hours. After evaporation of the methanol, the residue was taken up in water, the pH adjusted 5 to 4.8 with 2N HCl and extracted with dichloromethane to give 2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoic acid after evaporation as a foam (0.522 g ; 85%).

¹H NMR (CDCl₃, 400 MHz) δ : 4-4.25 (3H, m) ; 4.4-4.7 (2H, m) ; 6.9-7.5 (12H, m) ; 7.8 (2H, m).

10

Example 2

(2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid

A solution of methyl (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoylamino}-4-methylsulfanyl butyrate (0.135 g ; 0.23 mmol) in methanol (10 ml) was treated with 2N aqueous sodium hydroxide solution (0.24 ml) and heated at 60°C for 4 hours. After evaporation of the methanol, the aqueous reaction mixture was acidified to pH 7.5 with 6N HCl and purified on reverse phase silica eluting with a gradient of 50-60% methanol / ammonium carbonate buffer (2 g/l, pH 7). The appropriate 20 fractions were concentrated and freeze-dried to give the title compound as a solid.

Yield : 65%

¹H NMR (DMSO d₆, 400 MHz) δ : 1.7-1.9 (2H, m) ; 2 (3H, s) ; 2.15-2.35 (2H, m) ; 4.2-4.5 (5H, m) ; 4.8 (1H, m) ; 6.83 (1H, m) ; 7.1-7.6 (14H, m) ; 8.4 (1H, s).

Anal. Calculated for C₃₀H₂₉F₂N₃O₄S, H₂O

25	C 61.74	H 5.35	N 7.2	S 5.49
	Found	C 61.97	H 5.26	N 7.14
				S 5.05

MS (ESI) m/z 566 (MH⁺)

Example 3**Tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethylbenzoylamino]-4-methylsulfanylbutyrate**

Tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethylbenzoylamino]-4-methylsulfanylbutyrate was prepared by a similar method to that used for Example 1, but using tert-butyl (2S)-2-amino-4-methylsulfanylbutanoate (L-methionine tert-butyl ester) in place of L-methionine methyl ester.

The compound was purified by flash column chromatography dichloromethane/ethanol, (98:2) and further purified on reverse phase silica eluting with a gradient of 60-80% methanol / ammonium carbonate buffer (2 g/l, pH 7).

Yield : 52%

¹H NMR (CDCl₃, 400 MHz) δ : 1.42 (9H, s) ; 1.7-2 (2H, m) ; 2.15-2.25 (2H, m) ; 4-4.3 (3H, m) ; 4.45-4.6 (3H, m) ; 6 (1H, m) ; 6.8-7.6 (14H, m).

Anal. Calculated for C₃₄H₃₇F₂N₃O₄S

C 65.68 H 6.0 N 6.76 S 5.16

Found C 65.33 H 6.11 N 6.66 S 4.84

MS (ESI) m/z 622 (MH⁺)

Example 4**Methyl (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(thiazol-2-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate**

The title compound was prepared using a similar method to that used for Example 1, but using 1-(thiazol-2-yl)-2-(imidazol-1-yl)-ethanol in place of 1-(4-fluorophenyl)-2-(imidazol-1-yl)-ethanol.

Yield : 76%

¹H NMR (DMSO d₆ + AcOD) δ : 1.75-2 (2H, m) ; 2 (3H, s) ; 2.15-2.35 (2H, m) ; 3.64 (3H, s) ; 4.37 (1H, m) ; 4.65-4.9 (4H, m) ; 5.37 (1H, m) ; 7.15-7.95 (11H, m) ; 8.62 (1H, d) ; 9.09 (1H, s).

Anal. Calculated for C₂₈H₂₉FN₄O₄S₂, 0.8 H₂O, 1HCl

C 54.28 H 5.14 N 9.04 S 10.35

- 39 -

Found C 54.30 H 5.16 N 9.30 S 10.24
MS (ESI) m/z 569 (MH⁺)

Example 5

5 **(2S)-2-{2-(4-Fluorophenyl)-4-[2-(imidazol-1-yl)-1-(thiazol-2-yl)ethoxymethyl]benzoylamino}-4-methylsulfanyl butyric acid**

The title compound was prepared by a similar method to that used for Example 2.

Yield : 78%

¹H NMR (DMSO d₆ + CF₃COOD 400 MHz) δ : 1.8-2 (2H, m) ; 2 (3H, s) ; 2.2-2.4 (2H, m) ;
10 4.3 (1H, m) ; 4.6-4.9 (4H, m) ; 5.35 (1H, m) ; 7.1-7.4 (6H, m) ; 7.5-7.7 (2H, m) ; 7.8 (1H, m) ;
7.9 (1H, m) ; 9.09 (1H, s).

Anal. Calculated for C₂₇H₂₇FN₄O₄S₂, 0.5 H₂O

C 57.53 H 5.01 N 9.94 S 11.38

Found C 57.15 H 4.95 N 9.98 S 10.73

15 MS (ESI) m/z 555 (MH⁺)

Example 6

Methyl (2S)-2-{2-(4-fluorophenethyl)-5-[2-(imidazol-1-yl)-1-(4-fluorophenyl)-ethoxymethyl]benzoylamino}-4-methylsulfanyl butyrate

20 The title compound was prepared by a similar method to that used for Example 1, but using methyl 5-bromomethyl-2-(4-fluorophenethyl) benzoate in place of methyl 4-bromomethyl-2-(4-fluorophenyl)benzoate

Yield : 71%

¹H NMR (DMSO d₆ + CF₃COOD, 400 MHz) δ : 1.9-2.15 (2H, m) ; 2.05 (3H, s) ; 2.4-2.7
25 (2H, m) ; 2.75-3.1 (4H, m) ; 3.66 (3H, m) ; 4.25-4.7 (5H, m) ; 4.95 (1H, m) ; 7-7.30 (9H, m) ;
7.3-7.75 (4H, m) ; 9.04 (1H, s).

Anal. Calculated for C₃₃H₃₅F₂N₃O₄S, 0.5 H₂O, 1HCl

C 60.68 H 5.71 N 6.43 S 4.91

Found C 60.29 H 5.88 N 6.37 S 4.58

30 MS (ESI) m/z 608 (MH⁺)

The starting material was prepared as follows:

A mixture of dimethyl 4-bromoisophthalate (54.75g., 200.5 mmol), water (330 ml), tributylamine (55.63g., 300.7 mmol), 4-fluorostyrene(55.63g., 300.7 mmol) and bis(triphenylphosphine)palladium(II)chloride (2.81g., 4.01 mmol) was heated at reflux with stirring under an inert atmosphere for 6 hours. The reaction was cooled to ambient temperature and acidified to pH 2 with 2M HCl (700ml). The aqueous layer was removed and the residual solid washed with water (2L), dissolved in dichloromethane (1L) and passed through a pad of silica, eluting with more dichloromethane(2L). Evaporation of the dichloromethane gave a solid, which was washed with iso-hexane(1L) and dried to give methyl 4-[2-(4-fluorophenyl)ethenyl]-3-methoxycarbonylbenzoate (56.75g) as a pale yellow solid.

NMR data (CDCl₃) δ : 3.96 (6H, 2s), 7.01-7.10 (3H, m), 7.49-7.57 (2H, m), 7.80 (1H, d), 7.97 (1H, d), 8.16 (1H, dd), 8.60 (1H, s).

MS m/e 315.3 (M+H)⁺.

A mixture of methyl 4-[2-(4-fluorophenyl)ethenyl]-3-methoxycarbonylbenzoate (56.75g, 180.6 mmol), ethyl acetate (900 ml), 10 % palladium on carbon(6g) was stirred under an hydrogen atmosphere for 6 hours. The catalyst was filtered and replaced with fresh catalyst (6g). The reaction was then stirred under an hydrogen atmosphere for 16 hours. The catalyst was filtered and the filtrate evaporated to dryness to give, as a colourless gum, methyl 4-(4-fluorophenethyl)-3-methoxycarbonylbenzoate (55.00g).

NMR data (CDCl₃) δ : 2.84-2.93 (2H, m), 3.25-3.33 (2H, m), 3.93 (6H, 2s), 6.90-7.00 (2H, m), 7.09-7.16 (2H, m), 7.22-7.28 (1H, m), 8.05 (1H, dd), 8.57 (1H, s).

MS m/e 317.3 (M+H)⁺.

A mixture of methyl 4-(4-fluorophenethyl)-3-methoxycarbonylbenzoate (51.00g., 161.22 mmol), dioxane(650ml), methanol (650ml), sodium hydroxide (7.10g., 177.35 mmol) and water (100ml) was stirred at ambient temperature under an inert atmosphere for 16 hours. The reaction was evaporated to dryness, water (500 ml) added to the residue and the mixture extracted with diethylether. The organic extracts were dried and evaporated to dryness to give recovered methyl 4-(4-fluorophenethyl)-3-methoxycarbonylbenzoate (7 g). The aqueous layer was acidified to pH 2 with 2M HCl (300 ml) and extracted with ethyl acetate(300 ml). The organic extracts were dried, filtered and evaporated to dryness to give, as a white solid, 4-(4-fluorophenethyl)-3-methoxycarbonylbenzoic acid (42.00g).

NMR data (DMSO d_6) δ : 2.77-2.85 (2H, m), 3.16-3.24 (2H, m), 3.85 (3H, s),
7.04-7.12 (2H, m), 7.17-7.25 (2H, m), 7.45 (1H, d), 8.00 (1H, dd), 8.535 (1H, s).
MS m/e 301.4 (M-H)⁻.

A mixture of 4-(4-fluorophenethyl)-3-methoxycarbonylbenzoic acid (16.5g, 54.88
5 mmol), tetrahydrofuran (500 ml) and borane in tetrahydrofuran (1M. complex, 218mls, 218
mmol) was stirred under an inert atmosphere at reflux for 6 hours. The reaction was cooled to
ambient temperature and methanol (1L) was added. It was then evaporated to dryness to give
a dark oil which was purified by flash column chromatography using iso-hexane/ethyl acetate
(1:1) as eluant to give, as a clear gum, methyl 2-(4-fluorophenethyl)-5-
10 hydroxymethylbenzoate

(13.10g).

NMR data (CDCl₃) δ : 1.74 (1H, t), 2.82-2.92 (2H, m), 3.17-3.27 (2H, m), 3.91 (3H, s), 4.71
(2H, d), 6.91-6.99 (2H, m), 7.11-7.20 (3H, m), 7.41 (1H, d), 7.91 (1H, s).
MS m/e 289 (M+H)⁺.

15 A mixture of methyl 2-(4-fluorophenethyl)-5-hydroxymethylbenzoate (13.10g, 45.43
mmol), carbon tetrabromide (18.08g, 54.52 mmol) and triphenylphosphine (14.30g, 54.52
mmol) in dichloromethane (400 ml) was stirred at ambient temperature for 4 hours. More
carbon tetrabromide (7.54g, 23.00 mmol) and triphenylphosphine (5.96g, 23.00 mmol) in
dichloromethane (50 ml) were added. The reaction was applied directly to a silica flash
20 column and eluted with iso-hexane/ethyl acetate (92.5:7.5) to give methyl 2-(4-
fluorophenethyl)-5-bromomethylbenzoate (9.30g) as a clear gum.

NMR data (CDCl₃) δ : 2.81-2.91 (2H, m), 3.19-3.27 (2H, m), 3.91 (3H, s), 4.48 (2H, s), 6.91-
7.00 (2H, m), 7.12-7.17 (3H, m), 7.44 (1H, dd), 7.95 (1H, s).

MS m/e 's 351 and 353 (M+H)⁺.

25

Example 7

(2S)-2-{2-(4-fluorophenethyl)-5-[2-(imidazol-1-yl)-1-(4- fluorophenyl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid

The title compound was prepared by a similar method to that used for Example 2.

30 Yield : 73%

¹H NMR (DMSO d₆ + CF₃COOD, 400 MHz) δ : 1.9-2.15 (2H, m) ; 2.04 (3H, s) ; 2.5-2.7 (2H, m) ; 2.75-3.1 (4H, m) ; 4.2-4.6 (5H, m) ; 4.91 (1H, m) ; 7-7.75 (13H, m) ; 9.03 (1H, s).

Anal. Calculated for C₃₂H₃₃F₂N₃O₄S, H₂O

5	C 62.83	H 5.77	N 6.87	S 5.24	
	Found	C 62.48	H 5.66	N 6.87	S 5.06
	MS (ESI) m/z 594 (MH ⁺)				

Example 8

10 Methyl (2S)-2-{2-(4-(fluorobenzyl)-5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate

The title compound was prepared by a similar method to that used for Example 1, but using methyl 5-bromomethyl-2-(4-fluorobenzyl)benzoate in place of methyl 4-bromomethyl-2-(4-fluorophenyl)benzoate.

15 Yield : 48%

¹H NMR (CDCl₃, 400 MHz) δ : 2-2.25 (2H, m) ; 2 (3H, s) ; 2.45-2.65 (2H, m) ; 3.78 (3H, s) ; 4.1-4.9 (8H, m) ; 6.9-7.45 (14H, m) ; 9 (1H, d).

Anal. Calculated for C₃₂H₃₃F₂N₃O₄S, 0.5 H₂O, 1HCl

	C 60.13	H 5.52	N 6.57	S 5.02	
20	Found	C 60.00	H 5.57	N 6.61	S 4.87
	MS (ESI) m/z 594 (MH ⁺)				

The starting material was prepared as follows:

A 2.0M solution of 4-fluorobenzyl zinc bromide in tetrahydrofuran (95ml, 190mmol; Negishi, E-i. and King, A.O., J.O.C.1977,42,1821) was added dropwise over 20 minutes to a
 25 stirred solution of dimethyl 4-bromoisophthalate (20.0g, 73.2mmol) and tris(dibenzylideneacetone)dipalladium(0) (0.67g,0.73mmol) in tetrahydrofuran (270ml) at ambient temperature under an inert atmosphere. During the addition the internal temperature rose to 40°C. The reaction mixture was stirred for 3 hours and then poured into 2M. HCl (1500 ml) cooled in an ice bath and the product extracted into ethyl acetate (3x700ml). The
 30 extracts were washed with water (500ml), brine (500ml), dried, and the solvent evaporated under reduced pressure. The residue was taken up in a small quantity of dichloromethane,

isohexane added, and the solid, which crystallised from the solution, filtered and dried to give methyl 4-(4-fluorobenzyl)-3-methoxycarbonylbenzoate (10.5 g)

NMR data (CDCl₃) δ : 3.87 (3H, s), 3.95 (3H, s), 4.40 (2H, s), 6.93-7.00 (2H, m), 7.06-7.15 (2H, m), 7.29 (1H, d), 8.07 (1H, dd), 8.58 (1H, d).

5 MS m/e 303 (M+H)⁺

Methyl 4-(4-fluorobenzyl)-3-methoxycarbonylbenzoate (16.9g, 56.0 mmol) in methanol (270ml) was warmed slightly to achieve solution. 2.5M Sodium hydroxide (27ml, 67.5mmol) was added and the reaction mixture stirred at ambient temperature for 5 hours. The reaction mixture was poured into water (1200ml), acidified with concentrated HCl
10 and the product extracted into dichloromethane (3x400ml). The combined extracts were washed with brine, dried and the solvent evaporated under reduced pressure to give a mixture of 4-(4-fluorobenzyl)-3-methoxycarbonylbenzoic acid and the isomeric monoester, as a cream solid (15.8g).

NMR data (CDCl₃) δ : 3.85-3.98 (3H, m), 4.40-4.52 (2H, m), 6.94-7.02 (2H, m), 7.07-
15 7.17(2H, m), 7.32-7.38 (1H, m), 8.15-8.19 (1H, m), 8.63-8.75 (1H, m)

MS m/e 287 (M-H)⁻

A 1.0M solution of borane in tetrahydrofuran (100ml, 100 mmol) was slowly added to a stirred solution of the monoester mixture (15.8g, 54.7 mmol) at ambient temperature under an inert atmosphere. After the addition the reaction mixture was refluxed for 2 hours, cooled
20 to ambient temperature and methanol (100ml) added. The solvent was evaporated under reduced pressure and the product purified by column chromatography eluting with ethyl acetate/isohexane (35:65 and 40:60) to give methyl 5-hydroxymethyl-2-(4-fluorobenzyl)benzoate as a light yellow oil (9.3g).

NMR data (CDCl₃) δ : 1.82 (1H, t), 3.82 (3H, s), 4.35 (2H, s); 4.68-4.74 (2H, m), 6.90-6.99
25 (2H, m), 7.05-7.13 (2H, m), 7.21 (1H, d), 7.46 (1H, dd), 7.90 (1H, d)

MS m/e 275 (M+H)⁺

Triphenylphosphine (9.7g., 37.0 mmol) and then carbon tetrabromide (12.3g., 37.0 mmol) were added to a stirred solution of methyl 5-hydroxymethyl-2-(4-fluorobenzyl)benzoate (4.6g., 16.8mmol) in anhydrous ether (150ml) at ambient temperature.
30 After 4 hours the reaction was filtered and the filtrate concentrated under reduced pressure.

The product was purified by column chromatography eluting with ethyl acetate/isohexane (5:95) to give methyl 5-bromomethyl-2-(4-fluorobenzyl)benzoate as a colourless oil (5.05g). NMR data (CDCl₃) δ : 3.83 (3H, s), 4.37 (2H, s), 4.50 (2H, s), 6.92-6.99 (2H, m), 7.06 7.13 (2H, m), 7.19 (1H, d), 7.43 (1H, dd), 7.95 (1H, d).

5

Example 9**(2S)-2-{2-(4-fluorobenzyl)-5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid**

The title compound was prepared by a similar method to that used for Example 2.

10 Yield : 77%

¹H NMR (DMSO d₆ + CF₃COOD, 400 MHz) δ : 1.9-2.15 (2H, m) ; 2.04 (3H, s) ; 2.4-2.7 (2H, m) ; 4-4.75 (7H, m) ; 4.9 (1H, m) ; 7-7.7 (13H, m) ; 9.02 (1H, s).

Anal. Calculated for C₃₁H₃₁F₂N₃O₄S, 1.5 H₂O

	C 61.37	H 5.65	N 6.93	S 5.29	
15	Found	C 61.31	H 5.72	N 7.17	S 5.09

MS (ESI) m/z 580 (MH⁺).

Example 10**Methyl (2S)-2-{2-phenyl-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate**

20

The title compound was prepared by a similar method to that used for Example 1, but using methyl 4-bromomethyl-2-phenylbenzoate in place of methyl 4-bromomethyl-2-(4-fluorophenyl)benzoate.

Yield : 81%

25 ¹H NMR (DMSO d₆ + CF₃COOD, 400 MHz) δ : 1.75-2 (2H, m) ; 2 (3H, s) ; 2.15-2.30 (2H, m) ; 3.64 (3H, s) ; 4.30-4.60 (5H, m) ; 4.94 (1H, m) ; 7.1-7.8 (14H, m) ; 9.03 (1H, s).

Anal. Calculated for C₃₁H₃₂FN₃O₄S, 5.1 H₂O

	C 62.25	H 5.56	N 7.03	S 5.36	
30	Found	C 61.90	H 5.67	N 7.05	S 5.211

MS (ESI) m/z 562 (MH⁺)

The starting material was prepared by a similar method to that used for 4-bromomethyl-2-(4-fluorophenyl)benzoate in Example 1.

Example 11

5 (2S)-2-{2-phenyl-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid

The title compound was prepared by a similar method to that used for Example 2.

Yield : 90%

¹H NMR (DMSO d₆ + CF₃COOD, 400 MHz) δ : 1.8-2 (2H, m) ; 2 (3H, s) ; 2.1-2.35 (2H, m);
10 4.25-4.60 (5H, m) ; 4.9 (1H, m) ; 7.1-7.8 (14H, m) ; 9.07 (1H, s).

Anal. Calculated for C₃₀H₃₀FN₃O₉S, 0.12 H₂O

C 65.54	H 5.54	N 7.64	S 5.83
---------	--------	--------	--------

Found	C 65.13	H 5.47	N 7.71 S 5.86
-------	---------	--------	---------------

MS (ESI) m/z 548 (MH⁺).

15

Example 12

Methyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate

A solution of methyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate (compound of Example 1) (0.604
20 g ; 1.04 mmol) and 70 % MCPBA (0.650 g ; 2.6 mmol) in dichloromethane (15 ml) was stirred at ambient temperature for 5 hours. The solution was washed with saturated sodium bicarbonate and saturated sodium chloride solution and evaporated to dryness. The residue was purified on reverse phase silica eluting with a gradient of 60-70 % methanol /ammonium
25 carbonate buffer (2 g/l ; pH7). The appropriate fractions were concentrated and freeze-dried to give the title compound.

Yield : 63 %

The hydrochloride salt was prepared by addition of 3.86N HCl in ether (180 µl) to a solution of the above compound (0.32 g ; 0.52 mmol) in dichloromethane (3 ml). The mixture
30 was diluted with ether (100 ml) and the resulting precipitate was filtered and dried.

¹H NMR (DMSO-d₆ + CF₃COOD, 400 MHz) δ : 2-2.3 (2H, m) ; 2.9-3.2 (2H, m) ; 3.02 (3H, s) ; 3.72 (3H, s) ; 4.4-4.7 (5H, m) ; 5 (1H, m) ; 7.15-7.85 (13H, m) ; 8.85 (1H, d) ; 9.15 (1H, s).

Anal. Calculated for C₃₁H₃₁F₂N₃O₆S, 0.9 HCl, 0.6 H₂O

5 C 56.82 H 5.09 N 6.41 S 4.99 Cl 4.87
 Found C 57.27 H 5.55 N 6.03 S 4.57 Cl 5.07
 MS (ESI) m/z : 612 (MH⁺)

Example 13

10 (2S)-2-{2-(4-Fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyric acid

The title compound was prepared from Example 12 by a similar method to that used for Example 2.

Yield : 84 %

15 ¹H NMR (DMSO-d₆ + CF₃COOD, 400 MHz) δ : 1.9-2.3 (2H, m) ; 2.8-3.2 (2H, m) ; 2.96 (3H, s) ; 4.3-4.6 (5H, m) ; 4.96 (1H, m) ; 7.1-7.8 (13H, m) ; 8.8 (1H, d) ; 9.08 (1H, s).

Anal. Calculated for C₃₀H₂₉F₂N₃O₆S, 1.2 H₂O

 C 58.19 H 5.11 N 6.79 S 5.18
 Found C 58.22 H 5.24 N 6.78 S 4.93
 20 MS (ESI) m/z : 598 (MH⁺)

Example 14

tert-Butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate

25 The title compound was prepared from Example 3 by a similar method to that used for Example 12.

Yield : 67 %

¹H NMR (CDCl₃ + CD₃COOD, 400 MHz) δ : 1.43 (9H, s) ; 2-2.4 (2H, m) ; 2.7-3 (2H, m) ; 2.90 (3H, s) ; 4.2-4.8 (6H, m) ; 6.95-7.4 (12H, m) ; 7.56 (1H, m) ; 8.34 (1H, s).

30 Anal. Calculated for C₃₄H₃₇F₂N₃O₆S

 C 62.47 H 5.7 N 6.43 S 4.9

- 47 -

Found C 62.31 H 6.13 N 6.26 S 4.46

MS (ESI) m/z : 654 (MH⁺).**Example 15****5 N-Methylpiperidin-4-yl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate**

The title compound was prepared from 2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoic acid by a similar method to that used for Example 1 but using N-methylpiperidin-4-yl (2S)-2-amino-4-(methylsulfanyl)butanoate (L-methionine (N-methylpiperidin-4-yl)ester) in place of L-methionine methyl ester.

Yield : 72.5 %

¹H NMR (CDCl₃, 400 MHz) δ : 1.6-2.1 (6H, m) ; 2.03 (3H, s) ; 2.1-2.70 (6H, m) ; 2.27 (3H, s) ; 4.05-4.30 (3H, m) ; 4.45-4.80 (4H, m) ; 6.05 (1H, m) ; 6.8-7.6 (14H, m).

Anal. Calculated for

15 C 65.24 H 6.08 N 8.45 S 4.84

Found C 64.89 H 6.33 N 8.38 S 4.54

MS (ESI) m/z : 663 (MH⁺)**Example 16****20 Tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate**

The title compound was prepared from 2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxymethyl]benzoic acid by a similar method to that used for Example 1 but using L-methionine tert-butyl ester in place of L-methionine methyl ester.

25 Yield : 63 %

MP : 50 - 53°C

¹H NMR (CDCl₃, 400 MHz) δ : 1.42 (9H, s) ; 1.7-2.3 (4H, m) ; 2.04 (3H, s) ; 4.2-4.75 (5H, m) ; 5.01 (1H, m) ; 6.18 (1H, m) ; 6.9-7.5 (10H, m) ; 7.60 (1H, m) ; 7.82 (1H, m).

Anal. Calculated for C₃₁H₃₃FN₄O₄S₂

30 C 60.96 H 5.78 N 9.17 S 10.50

Found C 60.57 H 5.89 N 9.41 S 9.88

MS (ESI) m/z : 611 (MH^+).

Example 17

Tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate

The title compound was prepared from 2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxymethyl]benzoic acid by a similar method to that used for Example 1 but using the appropriate sulphonyl compound in place of the L-methionine methyl ester.

Yield : 76 %

10 MP : 90-100°C

1H NMR ($CDCl_3$, 400 MHz) δ : 1.43 (9H, s) ; 2-2.4 (2H, m) ; 2.6-2.9 (2H, m) ; 2.87 (3H, s) ; 4.20-4.60 (4H, m) ; 4.70 (1H, d) ; 4.99 (1H, m) ; 6.35 (1H, m) ; 6.85-7.45 (10H, m) ; 7.57 (1H, m) ; 7.84 (1H, m).

Anal. Calculated for $C_{31}H_{33}FN_4O_6S_2$

15 C 57.93 H 5.49 N 8.72 S 9.98

Found C 58.05 H 5.87 N 8.81 S 9.92

MS (ESI) m/z : 643 (MH^+).

Example 18

Methyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate

The title compound was prepared from Example 4 by a similar method to that used for Example 12.

Yield : 71 %

25 1H NMR ($CDCl_3$, 400 MHz) δ : 2-2.4 (2H, m) ; 2.7-3 (2H, m) ; 2.87 (3H, s) ; 3.71 (3H, s) ; 4.20-4.80 (5H, m) ; 5 (1H, m) ; 6.5 (1H, m) ; 6.93 (1H, d) ; 7.1-7.4 (9H, m) ; 7.56 (1H, m) ; 7.84 (1H, m).

Anal. Calculated for $C_{28}H_{29}FN_4O_6S_2, 0.4 H_2O$

C 55.32 H 4.94 N 9.22 S 10.55

30 Found C 55.07 H 5.11 N 8.88 S 10.24

MS (ESI) m/z : 601 (MH^+)

Example 19**(2S)-2-{2-(4-Fluorophenyl)-4-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyric acid**

The title compound was prepared from Example 18 by a similar method to that used
5 for Example 2, but heating at ambient temperature for 1 hour instead of at 60°C for 4 hours.

Yield : 85 %

¹H NMR (DMSO-d₆ + CF₃COOD, 400 MHz) δ : 1.95-2.3 (2H, m) ; 2.85-3.15 (2H, m) ;
2.96 (3H, s) ; 4.35 (1H, m) ; 4.6-4.90 (4H, m) ; 5.40 (1H, m) ; 7.15-8 (11H, m) ; 9.11 (1H, s).

Anal. Calculated for C₂₇H₂₇F N₄O₆S₂, 2 H₂O, 0.7 CH₃COONH₄

10 C 50.41 H 5.35 N 9.73 S 9.48

C 50.74 H 5.32 N 9.37 S 9.18

MS (ESI) m/z : 587 (MH⁺)

Example 20**15 N-Methylpiperidin-4-yl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate**

The title compound was prepared by a similar method to that used for Example 1 but
using L-methionine (N-methylpiperidin-4-yl) ester in place of L-methionine methyl ester.

Yield : 50 %

20 ¹H NMR CDCl₃, 400 MHz) δ : 1.65-2.1 (6H, m) ; 2.03 (3H, s) ; 2.1-2.7 (6H, m) ; 2.17
(3H, s) ; 4.25-4.80 (6H, m) ; 5.01 (1H, m) ; 6.25 (1H, m) ; 6.85-7.4 (10H, m) ; 7.6 (1H, m) ;
7.81 (1H, m).

Anal. Calculated for C₃₃H₃₈FN₅O₄S₂, 0.7 H₂O

C 59.65 H 5.98 N 10.54 S 9.65

25 Found C 59.72 H 5.90 N 10.37 S 9.16

MS (ESI) m/z : 652 (MH⁺).

Example 21**tert-Butyl (2S)-2-{2-(4-fluorophenethyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate**

The title compound was prepared from 2-(4-fluorophenethyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoic acid by a similar method to that used for Example 1 using the appropriate sulphonyl tert-butyl ester (tert-butyl (2S)-2-amino-4-(methylsulfonyl)butanoate) in place of the L-methionine methyl ester.

Yield : 30 %

¹H NMR (DMSO-d₆, 400 MHz) δ : 1.40 (9H, s) ; 2.1-2.4 (2H, m) ; 2.7-3.05 (4H, m) ; 3.08 (3H, s) ; 3.3-3.4 (2H, m) ; 4.20-4.50 (5H, m) ; 6.85 (1H, s) ; 7-7.55 (14H, m) ; 8.75 (1H, m).

Anal. Calculated for C₃₆H₄₁F₂N₃O₆S, 0.3 H₂O

C 62.92 H 6.10 N 6.11 S 4.67

Found C 62.57 H 6.26 N 6.06 S 4.33

MS (ESI) m/z : 682 (MH⁺).

Example 22**Tert-butyl (2S)-2-{2-(4-fluorophenethyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate**

The title compound was prepared from 2-(4-fluorophenethyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoic acid by a similar method to that used for Example 3.

Yield : 31 %

¹H NMR (CDCl₃, 400 MHz) δ : 1.50 (9H, s) ; 2.10 (3H, s) ; 2.05-2.35 (2H, m) ; 2.64 (2H, m) ; 2.85-3.15 (4H, m) ; 4-4.15 (3H, m) ; 4.4-4.55 (2H, m) ; 4.80 (1H, m) ; 6.8-7.45 (14H,).

Anal. Calculated for C₃₆H₄₁F₂N₃O₄S, 0.3 H₂O

C 65.99 H 6.40 N 6.41 S 4.89

Found C 65.99 H 6.78 N 6.41 S 4.55

MS (ESI) m/z : 650 (MH⁺).

Example 23**(2S)-2-{2-(4-Fluorophenethyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyric acid**

A solution of Example 21 (0.1 g ; 0.15 mmol) in solution in dichloromethane (0.5 ml) and TFA (1 ml) was stirred at ambient temperature for 2 hours. After evaporation to dryness, the residue was purified on reverse phase silica eluting with a gradient of 50-60 % methanol / ammonium carbonate buffer (2 g/l, pH 7). The appropriate fractions were concentrated and freeze dried to give the title compound.

Yield : 45 %

¹H NMR (DMSO-d₆ + CF₃COOD, 400 MHz) δ : 2.1-2.4 (2H, m) ; 2.75-3.1 (2H, m) ; 2.99 (3H, s) ; 3.1-3.4 (4H, m) ; 4.25-4.60 (5H, m) ; 4.90 (1H, m) ; 7-7.75 (13 H, m) ; 9.03 (1H, s).

Anal. Calculated for C₃₂H₃₃F₂N₃O₆S, 2 H₂O

C 58.08 H 5.64 N 6.35 S 4.85

Found C 57.88 H 5.30 N 6.27 S 4.97

MS (ESI) m/z : 626 (MH⁺).

Example 24**Tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(2-hydroxymethylimidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate**

The title compound was prepared by a similar method to that used for Example 1 but using L-methionine-t-butyl ester in place of L-methionine methyl ester

Yield : 60 %

¹H NMR (DMSO, 400 MHz) δ : 1.40 (9H, s) ; 1.7-2.3 (4H, m) ; 1.99 (3H, s) ; 4.15-4.50 (7H, m) ; 4.85 (1H, m) ; 5.26 (1H, m) ; 6.74 (1H, m) ; 7.05-7.5 (12H, m) ; 8.52 (1H, d).

Anal. Calculated for C₃₅H₃₉F₂N₃O₅S

C 64.5 H 6.03 N 6.45 S 4.92

Found C 64.74 H 6.05 N 6.23 S 4.57

MS (ESI) : 652 (MH⁺).

The starting material was prepared as follows :

A solution of 2-hydroxymethylimidazole (1.96 g ; 0.02 mole), imidazole (3.4 g ; 0.05 mole), tert-butyldiphenylsilyl chloride (6.6 g ; 0.024 mole) in DMF (20 ml) was stirred under an argon atmosphere for 16 hours. The mixture was extracted with ethyl acetate and purified by flash column chromatography eluting with a gradient 50-70 % ethyl acetate / petroleum ether to give 2-(tert-butyldiphenylsilyloxymethyl)imidazole as a solid.

Yield : 82 %

^1H NMR (CDCl_3 , 400 MHz) δ : 1.1 (9H, s) ; 4.85 (2H, s) ; 6.99 (2H, s) ; 7.3-7.5 (6H, m) ; 7.65 (4H, m).

A solution of 2-(tert-butyldiphenylsilyloxymethyl)imidazole (3.37 g ; 0.01 mol), bromomethyl acetone (2.17 g ; 0.01 mole) and TEA (1.67 ml ; 0.012 mole) in ethyl acetate (40 ml) was heated at reflux under argon atmosphere for 6 hours. After filtration of the insoluble material and evaporation to dryness, the residue was purified by flash column chromatography eluting with ethyl acetate / petroleum ether (50:50) to give 1-(4-fluorobenzoylmethyl)-2-(tert-butyldiphenylsilyloxymethyl)imidazole.

Yield : 71 %

^1H NMR (CDCl_3 , 400 MHz) δ : 0.96 (9H, s) ; 7.78 (2H, s) ; 5.53 (2H, s) ; 6.8-7.6 (14H, m) ; 7.95 (2H, m).

Sodium borohydride (0.41 g ; 1.06 mmol) was added portionwise at 0°C to a solution of 1-(4-fluorobenzoylmethyl)-2-(tert-butyldiphenylsilyloxymethyl)imidazole (4.81 g ; 1.01 mmol) in methanol (100 ml). The mixture was stirred at ambient temperature for 16 hours. 12N HCl (0.84 ml) added and then evaporated to dryness and purified by flash column chromatography eluting with dichloromethane / ethanol (96:4) to give 1-[2-(4-fluorophenyl)-2-hydroxyethyl]-2-(tert-butyldiphenylsilyloxymethyl)imidazole.

Yield : 63 %

^1H NMR (CDCl_3 , 400 MHz) δ : 1.10 (9H, s) ; 4.19 (2H, m) ; 4.61 (2H, s) ; 4.95 (1H, m) ; 6.8-7.8 (16H, m).

Methyl 2-(4-fluorophenyl)-4-{1-(4-fluorophenyl)-2-[2-(tert-butyldiphenylsilyloxymethyl)imidazol-1-yl]ethoxymethyl}benzoate was prepared by a similar method to that used for Example 1 but using 1-[2-(4-fluorophenyl)-2-hydroxyethyl]-2-(tert-butyldiphenylsilyloxymethyl)imidazole in place of 1-(4-fluorophenyl)-2-(imidazol-1-yl)ethanol

Yield : 90 %

¹H NMR (CDCl₃, 400 MHz) δ : 1.01 (9H, s) ; 3.65 (3H, s) ; 4.1-4.45 (4H, m) ; 4.60 (3H, m) ; 6.90-7.8 (23H, m).

Methyl 2-(4-fluorophenyl)-4-{1-(4-fluorophenyl)-2-[2-(tert-butyl)diphenylsilyloxymethyl]imidazol-1-yl]ethoxymethyl}benzoate (4 g ; 5.58 mmol) in methanol (50 ml) was treated with 2N aqueous sodium hydroxide solution (5 ml ; 11.1 mmol) at 80°C for 7 hours. After evaporation of the methanol, the residue was taken up in water, the pH adjusted to 5 with 6N HCl and extracted with dichloromethane / ethanol (95/5) to give 2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(2-hydroxymethylimidazol-1-yl)ethoxymethyl]benzoic acid.

Yield : 83 %

NMR (DMSO + CF₃COOD) δ : 4.2-4.4 (4H, m) ; 4.5-5 (3H, m) ; 7-7.7 (13 H, m).

15 **Example 25**

Methyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(2-hydroxymethylimidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate

The title compound was prepared from 2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(2-hydroxymethylimidazol-1-yl)ethoxymethyl]benzoic acid by a similar method to that used for Example 1.

Yield : 60 %

¹H NMR (CDCl₃, 400 MHz) δ : 1.7-2.3 (4H, m) ; 2.02 (3H, s) ; 3.68 (3H, s) ; 4.1-4.30 (3H, m) ; 4.45-4.8 (5H, m) ; 6.1-6.2 (1H, m) ; 6.84 (2H, m) ; 7.05-7.20 (6H, m) ; 7.22-7.4 (4H, m) ; 7.57 (1H, m).

25 Anal. Calculated for C₃₂H₃₃F₂N₃O₅S

C 63.04 H 5.46 N 6.89 S 5.26

Found C 63.40 H 5.64 N 6.64 S 4.84

MS (ESI) m/z : 610 (MH⁺)

Example 26**(2S)-2-{2-(4-Fluorophenyl)-4-[1-(4-fluorophenyl)-2-(2-hydroxymethylimidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid**

The title compound was prepared from Example 25 by a similar method to that used for Example 2.

Yield : 80 %

¹H NMR (CDCl₃ + CF₃COOD, 400 MHz) δ : 1.8-2.5 (4H, m) ; 2.06 (3H, s) ; 4.2-4.4 (3H, m) ; 4.5-5 (5H, m) ; 7-7.4 (12H, m) ; 7.6 (1H, m).

Anal. Calculated for C₃₁H₃₁F₂N₃O₅S, 0.5 H₂O

10	C 61.58	H 5.33	N 6.95	S 5.30
----	---------	--------	--------	--------

Found	C 61.63	H 5.55	N 6.85	S 4.91
-------	---------	--------	--------	--------

MS (ESI) m/z : 596 (MH⁺)

Example 27**15 Tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(2-hydroxymethylimidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate**

The title compound was prepared from 2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(2-hydroxymethylimidazol-1-yl)ethoxymethyl]benzoic acid and the appropriate tert-butyl ester by a similar method to that used for Example 1.

20 Yield : 42 %

¹H NMR (CDCl₃) : 1.43 (9H, s) ; 1.9-2.4 (2H, m) ; 2.6-2.95 (2H, m) ; 2.86 (3H, s) ; 4.1-4.3 (3H, m) ; 4.45-4.7 (5H, m) ; 6.15-6.25 (1H, m) ; 6.85 (2H, m) ; 7.05-7.4 (10H, m) ; 7.55 (1H, m).

Anal. Calculated for C₃₅H₃₉F₂N₃O₇S

25	C 61.48	H 5.75	N 6.15	S 4.69
----	---------	--------	--------	--------

Found	C 60.99	H 6.09	N 6.04	S 4.92
-------	---------	--------	--------	--------

MS (ESI) m/z : 684 (MH⁺)

Example 28**Methyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-2-methyl-4-methylsulfanylbutyrate**

The title compound was prepared by a similar method to that used for Example 1 but using α -methyl-L-methionine methyl ester.

Yield : 60 %

^1H NMR (CDCl_3 , 400 MHz) δ : 1.47 (3H, s) ; 1.9-2.6 (4H, m) ; 2.03 (3H, s) ; 3.70 (3H, s) ; 4.05-4.6 (5H, m) ; 6.34 (1H, s) ; 6.8-7.6 (14H, m).

Anal. Calculated for $\text{C}_{32}\text{H}_{33}\text{F}_2\text{N}_3\text{O}_4\text{S}$, 0.3 H_2O

10 C 64.16 H 5.65 N 7.01 S 5.35

Found C 63.95 H 5.65 N 6.84 S 4.80

MS (ESI) m/z : 594 (MH^+)

Example 29**(2S)-2-{2-(4-Fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-2-methyl-4-methylsulfanylbutyric acid**

The title compound was prepared by a similar method to that used for Example 2.

Yield : 42 %

^1H NMR (CDCl_3 , 400 MHz) δ : 1.57 (3H, d) ; 2.02 (3H, d) ; 2.05-2.3 (2H, m) ; 2.6-2.8 (2H, m) ; 4.05-4.25 (3H, m) ; 4.45-4.65 (2H, m) ; 6.8-7.35 (12H, m) ; 7.54 (1H, m) ; 7.86 (1H, s).

Anal. Calculated for $\text{C}_{31}\text{H}_{31}\text{F}_2\text{N}_3\text{O}_4\text{S}$, 0.6 H_2O

MS (ESI) m/z : 580 (MH^+)

25 Example 30**N-(4-Chlorobenzenesulfonyl)-(2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyramide**

The title compound was prepared by a similar method to that used for Example 1 but using N-(4-chlorobenzenesulfonyl)-(2S)-2-amino-4-methylsulfanylbutyramide.

30 Yield : 42 %

- 56 -

¹H NMR (CDCl₃ + CD₃COOD, 400 MHz) δ : 1.6-2.2 (4H, m) ; 1.95 (3H, d) ; 4.15-4.35 (2H, m) ; 4.45-4.70 (3H, m) ; 6.9-7.5 (15H, m) ; 7.90 (2H, m) ; 8.42 (1H, d).

Anal. Calculated for C₃₆H₃₃ClF₂N₄O₅S₂·0.3 H₂O

C 58.04 H 4.55 N 7.52 S 8.61

5 Found C 57.96 H 4.70 N 7.40 S 8.27

MS (ESI) m/z : 739 (MH⁺).

Example 31

2-(Morpholinomethyl)prop-2-yl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate

A mixture of 2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoic acid (0.462 g, 1 mmol), EDC (0.211 g, 1.1 mmol), 1-hydroxy-7-azabenzotriazole (0.148 g, 1.1 mmol), DMAP (0.122 g, 1 mmol) and 2-(morpholinomethyl)prop-2-yl (2S)-2-amino-4-methylsulfanylbutyrate (0.29 g, 1 mmol) in 15 dichloromethane (10 ml) was stirred at ambient temperature for 16 hours. The mixture was extracted with dichloromethane, evaporated to dryness and purified by flash column chromatography eluting with dichloromethane/ethanol (97:3) to give the title compound.

Yield : 60 %

¹H NMR (CDCl₃, 400 MHz) δ : 1.52 (6H, m) ; 2-2.34 (2H, m) ; 2.11 (3H, m) ; 2.51-2.79 (8H, m) ; 2.80-3.14 (4H, m) ; 3.67 (4H, m) ; 3.98-4.17 (3H, m) ; 4.4-4.53 (2H, m) ; 4.77 (1H, m) ; 6.81-7.48 (14H, m).

Anal. Calculated for C₄₀H₄₈F₂N₄O₅S

C 65.37 H 6.58 N 7.60 S 4.36

Found C 65.20 H 6.96 N 7.77 S 4.69

25 MS (ESI) m/z : 735 (MH⁺)

The starting material 2-(morpholinomethyl)prop-2-yl (2S)-2-amino-4-methylsulfanylbutyrate was prepared as follows.

A solution of N-(2-oxopropyl)morpholine (82 g, 0.57 mmol) in diethyl ether (400 ml) was added to a solution of 3M CH₃MgBr (500 ml, 1.5 ml) in anhydrous diethyl ether (2l) and 30 stirred at 0°C under an argon atmosphere. The mixture was stirred at ambient temperature for 16 hours, treated with 12N HCl (50 ml) and extracted with ethyl acetate. The product was

purified by flash column chromatography eluting with ethyl acetate to give N-(2-hydroxy-2-methylpropyl)morpholine.

Yield : 33 %

¹H NMR (CDCl₃, 400 MHz) δ : 1.05-1.4 (6H, m) ; 2.32 (2H, s) ; 2.55-2.72 (4H, m) ;
5 2.98 (1H, br) ; 3.6-3.82 (4H, m).

A solution of N-(2-hydroxy-2-methylpropyl)morpholine (9.8 g, 62 mmol), N-benzyloxycarbonyl-L-methionine (12.6 g, 44 mmol), bromo-tris-pyrrolidino-phosphonium hexafluorophosphate (25 g, 54 mmol), DMAP (5.4 g, 44 mmole) in dichloromethane (250 ml) was stirred, in the presence of 4 Å molecular sieves (5 g), at ambient temperature for 48 h.
10 After evaporation to dryness, the mixture was taken up in diethyl ether and the insoluble material eliminated by filtration. The organic layer was washed with a saturated aqueous solution of sodium hydrogen carbonate and a saturated solution of brine, evaporated to dryness and purified by flash column chromatography eluting with dichloromethane/ethanol (98:2) to give 2-(morpholinomethyl)prop-2-yl (2S)-2-(benzyloxycarbonylamino)-4-
15 methylsulfanylbutyrate (40 %).

¹H NMR (CDCl₃, 400 MHz) δ : 1.51 (6H, m) ; 1.85-2.27 (2H, m) ; 2.09 (3H, s) ; 2.45-2.70 (8H, m) ; 3.66 (4H, m) ; 4.36 (1H, m) ; 5.10 (2H, m) ; 5.25-5.40 (1H, m) ; 7.3-7.5 (5H, m).

A solution of HCO₂NH₄ (5.6 g, 88.5 mmol) in water (10 ml) was added dropwise to a
20 suspension of compound 2-(morpholinomethyl)prop-2-yl (2S)-2-(benzyloxycarbonylamino)-4-methylsulfanylbutyrate (7.5 g, 17.7 mmol) and 10% palladium on carbon (7.5g) in DMF (80 ml). The mixture was stirred at ambient temperature for 18 h. After filtration through Celite, the titrate was evaporated to dryness, neutralised with ammonia in diethyl ether to pH 9 and extracted with more ether. After evaporation the compound was purified by flash column
25 chromatography eluting with a gradient of 1-5% ethanol/dichloromethane to give 2-(morpholinomethyl)prop-2-yl (2S)-2-amino-4-methylsulfanylbutyrate (50 %).

¹H NMR (CDCl₃, 400 MHz) δ : 1.51 (6H, m) ; 1.6-2.08 (2H, m) ; 2.12 (3H, s) ; 2.50-2.80 (8H, m) ; 3.47 (1H, m) ; 3.62-3.77 (4H, m).

Example 32**Methyl (2S)-2-{4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenyl)benzoylamino}-4-methylsulfanylbutyrate**

ADDP (0.404 mg ; 1.53mmol) was added to a solution of 1-(4-fluorophenyl)-2-
5 (imidazol-1-yl) ethanol (0.3 g ; 1.45 mmol), methyl (2S)-2-[2-(4-fluorophenyl)-4-
hydroxybenzoylamino]-4-methylsulfanylbutyrate (0.49 g ; 1.45 mmol) and
triphenylphosphine (0.4 g ; 1.52 mmol) in THF (8 ml). The mixture was stirred for 16 hours
at ambient temperature. After evaporation to dryness, the residue was purified by flash
column chromatography eluting with a gradient of 1-20% ethanol / dichloromethane. The
10 product was further purified on reverse phase silica eluting with 60% methanol / ammonium
carbonate buffer (2 g/l, pH 7). The appropriate fractions were evaporated, redissolved in
dichloromethane (2 ml), treated at 0°C with a 3.8 N HCl solution in diethyl ether, diluted
with diethyl ether (100 ml) to precipitate the hydrochloride salt which was filtered and dried
to give the title compound (25%).

15 ¹H NMR (CDCl₃, 400 MHz) δ : 1.7-2.1 (2H, m) ; 2 (3H, s) ; 2.1-2.25 (2H, m) ; 3.65
(3H, s) ; 4.35 (2H, m) ; 4.60 (1H, m) ; 5.36 (1H, m) ; 5.82 (1H, d) ; 6.7-7.5 (14H, m).

MS (ESI) m/z 566 (MH⁺)

The starting material was prepared as follows:

Triethylamine (29 ml ; 0.206 mol) was added to a solution of methyl 4-
20 methoxysalicylate (25.0 g ; 0.137 mol) in dichloromethane (500 ml) and the solution cooled
to 0°C. Trifluoromethanesulphonic anhydride (29 ml ; 0.172 mol) was added dropwise and
the reaction stirred at ambient temperature for 1hour. Additional portions of triethylamine and
triflic anhydride were added over 16hours until HPLC showed absence of starting material.

The reaction was washed with 2N HCl and the organic phase evaporated to give a brown oil.
25 Purification by flash column chromatography eluting with ethyl acetate/iso-hexane (1:1) gave
methyl 4-methoxy-2-trifluoromethylsulfonyloxybenzoate as a pale yellow oil (23.4 g).

Yield : 76%

¹H NMR (CDCl₃, 300 MHz) δ : 3.88 (3H, s) ; 3.93 (3H, s) ; 6.79 (1H, d) ; 6.96
(1H, dd) ; 8.06 (1H, d).

30 MS (ESI) m/z 315 (MH⁺)

Saturated aqueous sodium hydrogen carbonate solution (50 ml) was added to a solution of methyl 4-methoxy-2-trifluoromethylsulfonyloxybenzoate (6.3 g ; 0.02 mol) and 4-fluorobenzeneboronic acid (3.36 g ; 0.024 mol) in DME (150 ml) at ambient temperature under an argon atmosphere. Tetrakis(triphenylphosphine) palladium (928 mg ; 0.8 mmol) was then added and the reaction heated at reflux for 3.5 hours to give a homogeneous solution. After cooling to ambient temperature, the reaction was partitioned between ethyl acetate and water. The organic phase was washed with 2N HCl, water and brine, filtered through 1PS filter paper and the solvent removed *in vacuo* to give methyl 4-methoxy-2-(4-fluorophenyl)benzoate as a yellow oily solid (7.2 g) which was used without further purification.

^1H NMR (CDCl_3 , 300 MHz) δ : 3.65 (3H, s) ; 3.87 (3H, s) ; 6.79 (1H, d) ; 6.91 (1H, dd) ; 7.08 (2H, dd), 7.25 (2H, dd), 7.90 (1H, d).

MS (ESI) m/z 261 (MH^+)

To a solution of methyl 4-methoxy-2-(4-fluorophenyl)benzoate (9.8 g ; 0.029 mol) in methanol (75 ml) was added 2N aqueous sodium hydroxide solution (45 ml) and the mixture heated at reflux for 1.5h. The reaction was cooled to ambient temperature, filtered and the filtrate concentrated *in vacuo* to remove the methanol. The residual aqueous phase was washed with diethyl ether, acidified to pH 1 with concentrated HCl and extracted with ethyl acetate. The organic extracts were dried (MgSO_4) and the solvent removed *in vacuo* to give 4-methoxy-2-(4-fluorophenyl)benzoic acid as a white solid (7.7 g), which was used without further purification.

^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ : 3.80 (3H, s) ; 6.80 (1H, d) ; 6.98 (1H, dd) ; 7.18 (2H, dd), 7.31 (2H, dd), 7.76 (1H, d).

MS (ESI) m/z 247 (MH^+)

A solution of boron tribromide (0.066mol) in dichloromethane (66 ml) was added dropwise to a solution of 4-methoxy-2-(4-fluorophenyl)benzoic acid (7.7 g ; 0.029 mol) in dried dichloromethane (215 ml) under argon at 0°C . The reaction was stirred for 1 hour at 0°C and allowed to warm to ambient temperature and stirred for a further 16 hours. It was then poured into ice water and extracted with firstly dichloromethane then with ethyl acetate. The combined organic extracts were washed with saturated aqueous sodium hydrogen carbonate and the aqueous phase acidified to pH 1 with concentrated HCl and extracted with ethyl

acetate. The ethyl acetate extracts were dried (MgSO_4) and concentrated *in vacuo* to give 4-hydroxy-2-(4-fluorophenyl)benzoic acid as a yellow oil (4.5 g), which was used without further purification.

^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ : 6.63 (1H, d) ; 6.80 (1H, dd) ; 7.15 (2H, dd), 7.26 (2H, dd), 7.71 (1H, d).

MS (ESI) m/z 233 (MH^+)

To a solution of 4-hydroxy-2-(4-fluorophenyl)benzoic acid (4.5 g ; 0.019 mol) in DMF (90 ml) cooled to 0°C under an argon atmosphere was added in sequence NMM (6.4 ml ; 0.058 mol), *L*-methionine methyl ester hydrochloride (4.0 g ; 0.020 mol) , EDC (4.47 g ; 0.023 mol) and HOBT (2.7 g ; 0.020 mol) and the reaction warmed to ambient temperature and stirred for 3 hours. The DMF was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic phase was washed with water and brine, filtered through PS1 filter paper and concentrated *in vacuo* to give a yellow oil. Purification by flash column chromatography eluting with ethyl acetate / *iso*-hexane (1:1) gave methyl (2S)-2-[4-hydroxy-2-(4-fluorophenyl)benzoylamino]-4-methylsulfanylbutyrate as a white foam (5.7 g).

Yield : 51% (over 4 steps)

^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ : 1.80 (1H, m) ; 1.95 (1H, m) ; 2.03 (3H, s) ; 2.20 (2H, dd) ; 3.70 (3H, s) ; 4.65 (1H, m) ; 5.97 (1H, d), 6.70 (1H, d) ; 6.78 (1H, dd) ; 6.82 (1H, br s) ; 7.07 (2H, dd), 7.30 (2H, dd), 7.55 (1H, d).

Anal. Calculated for $\text{C}_{19}\text{H}_{20}\text{FNO}_4\text{S}$

C 60.46	H 5.34	N 3.71	S 8.50	F 5.03
Found	C 60.1	H 5.4	N 3.6	S 8.3
	F 5.2			

MS (ESI) m/z 378 (MH^+)

Example 33

(2S)-2-{4-[1-(4-Fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenyl)benzoylamino}-4-methylsulfanylbutyric acid

A solution of methyl (2S)-2-{4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenyl)benzoylamino}-4-methylsulfanylbutyrate (0.037 g ; 0.061 mmol) in methanol (3 ml) was treated with 2N aqueous sodium hydroxide solution (0.32 ml) and the mixture stirred

at ambient temperature for 2 hours. After evaporation of the methanol, the aqueous residue was acidified to pH 7.5 with 6N HCl and purified on reverse phase silica eluting with a gradient of 50-60% methanol / ammonium carbonate buffer (2 g/l, pH 7). The appropriate fractions were concentrated and freeze-dried to give the title compound as a solid.

5 Yield : 42%

¹H NMR (DMSO d₆, 400 MHz) δ : 1.7-2 (2H, m) ; 2 (3H, s) ; 2.1-2.3 (2H, m) ; 4.16 (1H, s) ; 4.42 (2H, m) ; 5.8 (1H, m) ; 6.8-7.7 (14H, m) ; 8.25 (1H, m).

Anal. Calculated for C₂₉H₂₇F₂N₃O₄S, 0.9 H₂O

	C 61.34	H 5.11	N 7.4	S 5.65
10 Found	C 61.27	H 4.75	N 7.20	S 5.36

MS (ESI) m/z 552 (MH⁺)

Example 34

Methyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate

A mixture of 5-(1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy)-2-(4-fluorophenethyl)benzoic acid (570mg, 1.27 mmol), L-methionine methyl ester (510 mg, 2.5 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (370 mg, 1.9 mmol) in 20 dichloromethane (50ml), was stirred at ambient temperature for 16 hours. It was then washed with aqueous citric acid (2x40 ml, 1M), brine, dried (MgSO₄) and evaporated to dryness and the residue so obtained purified by chromatography on silica (40gm Dynamax column, gradient elution with 2-6% methanol /dichloromethane). The relevant fractions were combined and evaporated to give 640mg of a product which was re-dissolved in 25 dichloromethane, washed with aqueous citric acid (4x30 ml, 1M), brine, dried and evaporated to give a residue. This was redissolved in dichloromethane (4ml) at 0°C , treated with 1MHCl/Et₂O (10 ml), stirred for 2 hours, the solution evaporated to dryness and the residue triturated with ether to give the hydrochloride salt of the title compound as a white solid (343 mg).

30 ¹H NMR (CDCl₃, 300MHz) δ : 2-2.1 (4H,m) ; 2.15 - 2.32 (1H, m) ; 2.55 (2H,t); 2.72 -2.82 (2H,m) ; 2.84-2.98 (2H, m) ; 3.78 (3H,d) ; 4.2-4.4 (2H,m); 4.78 -4.86 (1H,m); 5.25 (-5.3

(1H,m); 6.34 (1H,d); 6.7 (1H,dd); 6.8-6.83 (1H,m); 6.84 - 7.00 (4H,m); 7.0 -7.12 (5H,m); 7.2-7.3 (2H,m); 7.44 (1H,s).

MS (ES+) m/z 594.3 (MH+)

Anal. calculated for $C_{32}H_{33}F_2N_3O_4S.HCl.H_2O$ C 59.3 H 5.6 N 6.5 S 4.9

5 Found C 59.3 H 5.5 N 6.1 S 5.2%

Example 35

(2S)-2-{5-[1-(4-Fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyric acid

10 A solution of methyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate (140 mg, 0.24 mmol) in methanol (5ml) was treated with sodium hydroxide (190mg,4.7 mmol) in water (1ml) and the mixture stirred at ambient temperature for 6 hours. The mixture was acidified to PH1 with 1M HCl and evaporated to dryness. The residue was washed with water (3 x 2mL) and the resulting

15 gum was triturated with diethyl ether to give the title compound as a white solid (43mg).
 1H NMR (DMSO- d_6 , 300MHz) δ : 1.85-2.05 (5H,m) ; 2.4-2.6 (signed partially obscured by solvent signals, m) ; 2.6-2.75 (2H,m); 2.75 -2.86 (2H,m) ; 4.4-4.48 (1H, m) ; 4.6-4.7 (2H,m) ; 5.82 (1H,t); 6.76-6.86 (2H,m); 6.96-7.08 (3H,m); 7.12-7.28 (4H,m); 7.44-7.5 (2H,dd); 7.62 (1H,s); 7.69 (1H,s); 8.58 (1H,t); 9.12 (1H,s).

20 MS (ES+) m/z 580.3 (MH+)

Anal. calculated for $C_{31}H_{31}F_2N_3O_4S.HCl.0.9NaCl$ C 55.7 H 4.8 N 6.3%

Found C 55.9 H 4.8 N 6.1%

Example 36

tert-Butyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate

The title compound was prepared in 80% yield from 5-(1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy)-2-(4-fluorophenethyl)benzoic acid and L-methionine tert-butyl ester by a similar method to that used for Example 34.

30 1H NMR (DMSO- d_6 , 300MHz) δ : 1.38 (9H,s) ; 1.88-2.0 (2H,m); 2.02 (3H,d); 2.62-2.75 (2H,m); 2.75 -2.9 (2H,m) ; 4.32-4.42 (1H, m) ; 4.68 (2H,d) ; 5.83 (1H,m); 6.79-6.83 (2H,md);

- 63 -

6.95-7.1 (3H,m); 7.1-7.28 (4H,m); 7.45-7.52 (2H,dd); 7.61 (1H,s); 7.68 (1H,s); 8.62 (1H,t); 9.08 (1H,s).

MS (ES+) m/z 636.4 (MH+)

Anal. calculated for $C_{35}H_{39}F_2N_3O_4S \cdot 2.7H_2O$ C 61.4 H 6.5 N 6.1%

5 Found C 61.1 H 6.2 N 5.7%

Example 37

Cyclopentyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate

10 The title compound was prepared in 72% yield from 5-(1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy)-2-(4-fluorophenethyl)benzoic acid and cyclopentyl (2S)-2-amino-4-(methylsulfanyl)butanoate (L-methionine cyclopentyl ester) by a similar method to that used for Example 34.

1H NMR (DMSO- d_6 , 300MHz) δ : 1.44-1.65 (6H,m) ; 1.7-1.86 (2H,m); 1.9-2.0 (2H,m);

15 2.02 (3H,d); 2.5 -2.6 (signals partially obscured by solvent signals,m); 2.62-2.9 (4H, m) ; 4.4-4.5 (1H,dd) ; 4.65 (2H,d); 5.04-5.1 (1H,m); 5.82 (1H,bds); 6.78-6.84 (2H,m); 6.97-7.1 (3H,dd); 7.12-7.3 (4H,m); 7.42-7.5 (2H,m); 7.58 (1H,bds); 7.74 (1H,bds); 8.68 (1H,t); 8.98 (1H,bds).

MS (ES+) m/z 648.4 (MH+)

20 Anal. calculated for $C_{36}H_{39}F_2N_3O_4S \cdot 2.8H_2O$ C 61.9 H 6.4 N 6.0

Found C 61.7 H 6.0 N 5.6

Example 38

tert-Butyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate

25 A mixture of 5-[(1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy)]-2-(4-fluorophenethyl)benzoic acid (800mg, 1.79 mmol), tert-butyl (2S)-2-amino-4-(methylsulfonyl)butanoate (L-methionine sulphone tert-butyl ester) (630mg, 2.68 mmol), and DMAP (870mg, 7.1 mmol), was treated with EDC.HCl (510 mg, 2.7 mmol) was stirred at
30 ambient temperature for 16 hours. The reaction mixture was washed with aqueous citirc acid (1M,2x), brine, dried ($MgSO_4$) and evaporated to dryness. The residue was purified by

chromatography on silica (40gm Dynamax column, gradient elution 2-10% methanol / dichloromethane) to give a gum which, when triturated with diethyl ether/iso-hexane, gave the title compound as a cream solid (889mg, 74%).

¹H NMR (DMSO-d₆, + HOAc-d₄, 300MHz) δ : 1.38 (9H,s) ; 2-2.25 (2H,m); 2.62-2.82 (4H,m); 2.97 (3H,d); 3.02-3.28 (2H,m); 4.36-4.42 (1H,dd) ; 4.6-4.67 (2H,d) ; 5.76-5.82 (1H,m) ; 6.78-6.84 (1H,m); 6.86-6.92 (1H,m); 6.92-7.05 (3H,m); 7.1-7.24 (4H,m); 7.4-7.58 (3H,m); 7.64-7.7 (1H,bds); 8.88-9.0 (1H,bds).

MS (ES+) m/z 668.3 (MH+)

Anal.calculated for C ₃₅ H ₃₉ F ₂ N ₃ O ₆ S.3H ₂ O	C	58.2	H	6.2	N	5.8
10 Found	C	58.1	H	6.0	N	5.6

The starting material was prepared as follows :

A mixture of methyl 2-bromo-5-methoxybenzoate (28.3 g, 115mmol), 4-fluorostyrene (20.6 ml, 173.3 mmol), tert-n-butylamine (2.7ml, 11.5 mmol), Pd[(o-tolyl)₃P]₂Cl₂ (0.9g, 1.15 mmol), and NaHCO₃ (14.6g, 173.3 mmol), and water (220ml) was stirred and heated at reflux for 16 hours. The reaction mixture was diluted with water (200ml) and extracted with ethyl acetate (2x400ml). The extracts were washed with 2M HCl (100ml), water (100ml), brine (100ml), dried (MgSO₄), filtered and evaporated to dryness. The brown gum obtained was purified by chromatography on silica (90g, Biotage column), eluting with a gradient of ethyl acetate in iso-hexane (0% - 20% ethyl acetate). The relevant fractions were combined and evaporated to give methyl 2-[2-(4-fluorophenyl)ethenyl]-5-methoxybenzoate, (43.5g)

¹H NMR (CDCl₃, 250MHz) δ : 3.86 (3H,s); 3.94 (3H,s); 6.86 (1H,d); 7.02 (1H,d); 7.04 - 7.12 (3H,m); 7.42 - 7.52 (3H,m); 7.64 (1H,d); 7.83 (1H,d).

MS (ES+) m/z 287 (MH+).

25 Methyl 2-[2-(4-fluorophenyl)ethenyl]-5-methoxybenzoate (43g, 150 mmol) in ethyl acetate (3.5L) was stirred with 10% Pd/C (4g) under an atmosphere of hydrogen (pressure 5 bar), at ambient temperature for 18 hours. The spent catalyst was removed by filtration through a Celite pad, the pad was washed with ethyl acetate and the filtrate evaporated to dryness to give methyl 2-(4-fluorophenethyl)-5-methoxybenzoate (33.8g, 80% yield).

30 ¹H NMR (CDCl₃, 300MHz) δ: 2.84 (2H, t); 3.16 (2H, t); 3.8 (3H, s); 3.88 (3H,s); 6.86-7.0 (3H,m); 7.06 (1H,d); 7.08-7.2 (2H, m); 7.43 (1H, d).

MS (ES+) m/z 289 (MH+).

Methyl 2-(4-fluorophenethyl)-5-methoxybenzoate (33.8g, 117 mmol) and pyridine hydrochloride (170g) were heated at 220°C for 2 hours. The black reaction mixture was cooled, poured into 2N HCl (1L), and the pale-brown solid extracted with ethyl acetate (2x 450ml). The extracts were washed with 2N HCl (500ml), brine (200ml), dried (MgSO₄) and evaporated to give 2-(4-fluorophenethyl)-5-hydroxy benzoic acid (30g, quantitative).

¹H NMR (DMSO-d₆, 300MHz) δ: 2.74 (2H, t); 3.04 (2H, t); 6.81 (1H, dd); 7.12 (3H, t); 7.15-7.24 (3H, m).

A solution of 2-(4-fluorophenethyl)-5-hydroxybenzoic acid (30g, 115.4 mmol) in methanol (900ml) and the stirred solution treated dropwise with sulphuryl chloride (1.6ml, 19.6 mmol) stirred and heated at reflux for 16 hours. The mixture was evaporated to dryness, the residue dissolved in ethyl acetate (500ml), washed with saturated aqueous NaHCO₃ (x2), water and brine. After drying (MgSO₄) the solution was evaporated to dryness to give a brown oil which when triturated with iso-hexane gave methyl 2-(4-fluorophenethyl)-5-hydroxybenzoate (21.02g, 66%).

¹H NMR (CDCl₃, 300MHz) δ: 2.84 (2H, t); 3.15 (2H, t); 4.8 - 5.2 (1H, broad); 7.85 - 7.0 (3H, m) 7.02 (1H, d); 7.12 (2H, dd); 7.39 (1H, d).

MS (ES+) m/z 275 (MH+).

Anal. calculated for C₁₆H₁₅F O₃

C 70.06 H 5.51

20 Found

C 70.0 H 5.6

A mixture of 1-(4-fluorophenyl)-2-imidazol-1-ylethanol, (5.0g, 24 mmol), methyl 2-(4-fluorophenethyl)-5-hydroxybenzoate, (7.3g, 27 mmol), DEAD (4.2 ml, 4.65g, 27 mmol), and triphenyl phosphine (7.0g, 27 mmol) in dry THF (150 ml) was stirred at ambient temperature for 16 hours. The reaction was evaporated to dryness and the residue purified by chromatography on silica (90g Biotage column, gradient elution 1-20% methanol/dichloromethane). The appropriate fractions were combined and evaporated to dryness to give methyl 5-[1-(4-fluorophenyl)-2-imidazol-1-ylethoxy]-2-(4-fluorophenethyl)benzoate (13.46g). The material was used without further purification in the next reaction.

30 MS (ES+) m/z 463 (MH+).

Methyl {5-[1-(4-fluorophenyl)-2-imidazol-1-yl]ethoxy}-2-(4-fluorophenethyl) benzoate (13.46g, ~29 mmol), was dissolved in methanol (100ml) and stirred at ambient temperature, with 2N aqueous sodium hydroxide solution (73ml, 146 mmol) for 4 days. The methanol was removed by evaporation and the residual aqueous solution adjusted to PH 7 with 2N HCl before being evaporated to dryness. The resulting white solid was purified by chromatography on silica (90g Biotage column, gradient elution 2-20%methanol/dichloromethane) to give 5-[1-(4-fluorophenyl)-2-imidazol-1-yl]ethoxy}-2-(4-fluorophenethyl)benzoic acid (4.0g), as a colourless foam.

¹H NMR (CDCl₃, 300MHz) δ: 2.84 (2H, t); 3.16 (2H, t); 4.34 (2H,d); 5.26-5.37 (1H, m); 6.66 (1H, dd); 6.88 (3H,t); 7.0 - 7.18 (5H,m); 7.22 - 7.35 (2H, m); 7.54 (1H, d); 7.92 (1H, s). MS (ES-) m/z447 (MH-).

Example 39

2-{5-[1-(4-Fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid

A solution of tert-butyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate, (88mg, 0.13mmol) in TFA (2ml), was stirred and cooled in an ice-bath for 3 hours. The TFA was removed by evaporation, the residue was dissolved in ethyl acetate and 1M HCl/diethylether (5ml). Removal of the solvent gave a brown oil which on trituration with diethylether/iso-hexane (3x) gave the HCl salt of the title compound as a sticky white solid (40mg).

¹H NMR (DMSO-d₆, 373°K, 250MHz) δ : 2.1-2.42 (2H,m) ; 2.75-2.85 (2H,m); 2.85-2.94 (2H,m); 2.95 (3H,d); 3.05-3.35 (2H,m); 4.5-4.6 (1H, m) ; 4.62-4.68 (2H,m) ; 5.73 (1H,t); 6.78-6.88 (1H,dd); 6.9-7.1 (4H,m); 7.1-7.25 (4H,m); 7.3-7.65 (4H,m); 8.09 (1H,d); 8.59 (1H,bds).

MS (ES+) m/z 612.3 (MH+)

Example 40**Methyl (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate**

A mixture of 5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoic acid (1.1g, 2.52 mmol), L-methionine methyl ester .HCl (1.0 g, 5.0 mmol), DMAP (1.54g, 12.6 mmol), in dichloromethane (40ml) and EDC.HCl (0.72 g 3.8mmol) was stirred for 16 hours. The mixture was washed with 1M citric acid (x2) and with brine, dried (MgSO₄), evaporated to dryness and the residue obtained purified by chromatography on silica (40gm Biotage column, gradient elution with 2 - 10% methanol/ 10 dichloromethane). The appropriate fractions were combined and evaporated to give the title compound (0.14gm, 64%) .

¹H NMR (DMSO-d₆, 300MHz) δ : 1.94-2.04 (5H,m) ; 2.52-2.6 (2H,m); 2.65-2.85 (4H,m); 3.6 (3H,s); 4.52 -4.65 (3H,m) ; 5.98-6.05 (1H, m) ; 6.82 (1H,s) ; 6.88-6.96 (2H,m); 7.0-7.2 (6H,m); 7.5 (1H,d); 7.76 (1H,d); 7.86 (1H,d); 8.64 (1H,t).

15 MS (ES+) m/z 583.3 (MH⁺)

Anal.calculated for C ₂₉ H ₃₁ FN ₄ O ₄ S ₂ .0.3H ₂ O	C	59.2	H	5.4	N	9.5
Found	C	59.2	H	5.3	N	9.3

The starting material was prepared as follows :

A mixture of 2-bromo-1-(thiazol-2-yl)-ethanone (2.5g; 12 mmol), acetonitrile (70ml) and imidazole (1.65g; 24 mmol) was stirred at ambient temperature for 1 hour. The mixture was evaporated to dryness and the residue partitioned between dichloromethane and water. The organic phase was separated, washed with saturated brine and evaporated to give 2-(imidazol-1-yl)-1-(thiazol-2-yl)-ethanone as a brown solid (1.17g; 50%).

MP : 109-112°C

25 ¹H NMR (DMSO-d₆, 400MHz) δ : 5.6 (2H,s) ; 6.85-8.1 (5H,m).

2-(Imidazol-1-yl)-1-(thiazol-2-yl)ethanone was reduced with sodium borohydride in methanol to give 2-(imidazol-1-yl)-1-(thiazol-2-yl)ethanol in 60% yield.

¹H NMR (CDCl₃, 300MHz) δ 4.15 (1H,dd); 4.57 (1H,dt); 5.2 (1H, dd); 6.75 (1H,s); 6.89 (1H,s); 7.32-7.41 (2H,m); 7.79 (1H,d).

30 MS (ES+) m/z 196 (MH⁺).

Methyl 5-[2-(imidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoate was prepared from 2-(imidazol-1-yl)-1-(thiazol-2-yl)ethanol and methyl 2-(4-fluorophenethyl)-5-hydroxybenzoate by the Mitsunobu methodology used for the preparation of methyl 5-[2-(2-methylimidazol-1-yl)-1-(4-fluorophenyl)ethoxy]-2-(4-fluorophenethyl)benzoate in Example 45.

¹H NMR (CDCl₃, 300MHz) δ 2.8 (2H,t); 3.12 (2H,t); 3.88 (3H,s); 4.52 (1H,dd); 4.62 (1H,dd); 5.76 (1H, dd); 6.85-6.97 (4H,m); 7.0-7.15 (4H,m); 7.34 (1H,d); 7.45 (1H,d); 7.48(1H,s); 7.86 (1H,d).

MS (ES+) m/z 452 (MH)+.

10 Methyl 5-[2-(imidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoate was hydrolysed using aqueous sodium hydroxide solution to give 5-[2-(imidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoic acid.

¹H NMR (CDCl₃, 300MHz) δ 2.85 (2H,t); 3.2 (2H,t); 4.54 (1H,dd); 4.68 (1H,dd); 5.82(1H,dd); 6.86-7.04 (5H,m); 7.04-7.16 (3H,m); 7.36 (1H,d); 7.59 (1H,d); 7.88 (2H,t).

15 MS (ES+) m/z 438 (MH)+.

Example 41

(2S)-2-{5-[1-(2-thiazolyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyric acid

20 Methyl (2S)-2-{5-[1-(2-thiazolyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate (140mg, 0.24 mmol) was dissolved in methanol (3ml), and treated with NaOH (190 mg, 4.8 mmol) in H₂O (1ml). After 4 hours at ambient temperature, the mixture was acidified to PH 1 with 1M HCl and the mixture evaporated to dryness. The residue was washed with water (3x2ml) and the resulting gum
25 triturated with iso-hexane to give the HCl salt of the title compound (94mg, 69%) as a white solid .

¹H NMR (DMSO-d₆, HOAc-d₄, 300MHz) δ : 1.84-2.1 (5H,m) ; 2.5-2.6 (signals partially obscured by the solvent signals, m); 2.65-2.78 (2H,m); 2.78-2.9 (2H,m); 4.4-4.51 (1H,dd) ; 4.8-4.98 (2H, m); 6.18-6.24 (1H,m); 6.8-7.04 (4H,m); 7.06-7.2 (3H,m); 7.59 (1H,s); 7.74 (1H,d); 7.86 (1H,d); 9.14 (1H,m).

MS (ES+) m/z 569.2 (MH+)

Anal. calculated for $C_{32}H_{37}FN_4O_6S_2 \cdot HCl \cdot 1.1H_2O \cdot 0.5NaCl$:

	C	51.4	H	5.0	N	8.6
Found	C	51.5	H	4.7	N	8.5

5 Example 42

tert-Butyl (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate

The title compound was prepared in 42% yield from 5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenylethyl)benzoic acid and L-methionine tert-butyl ester by a similar method to that used for Example 40.

1H NMR (DMSO- d_6 , 300MHz) δ : 1.39 (9H,s) ; 1.9-2.0 (2H,m); 2.02 (3H,s); 2.5-2.6 (2H,m); 2.68-2.78 (2H,m) ; 2.78-2.9 (2H, m) ; 4.36-4.45 (1H,m) ; 4.58-4.7 (2H,m); 6.0-6.05 (1H,m); 6.86-6.96 (2H,m); 6.98-7.08 (2H,t); 7.1-7.2 (4H,m); 7.5-7.65 (2H,m); 7.73 (1H,d); 7.86 (1H,d); 8.63 (1H,t).

MS (ES+) m/z 625.3 (MH+)

Anal. calculated for $C_{32}H_{37}FN_4O_4S_2 \cdot 1.25C_6H_8O_7$ (Citric acid)

	C	54.9	H	5.5	N	6.5
Found	C	54.9	H	5.5	N	6.7

20

Example 43

tert-Butyl (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate

The title compound was prepared in 75% yield from 5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoic acid and L-methionine sulphone tert-butyl ester by similar methodology to that used for Example 40.

1H NMR (DMSO- d_6 , HOAc- d_4 , 300MHz) δ : 1.37 (9H,s) ; 2.0-2.26 (2H,m); 2.6-2.8 (4H,dd); 2.75-2.94 (3H,m); 2.98 (3H,d) ; 3.1-3.38 (3H,m) ; 4.35-4.44 (1H,m) ; 4.7-4.84 (2H,m); 6.1-6.18 (1H,dd); 6.9-7.05 (3H,m); 7.08-7.2 (4H,m); 7.24 (1H,s); 7.48 (1H,s); 7.74 (1H,d); 7.88 (1H,d); 8.4 (1H,s); 8.72 (1H,t).

30 MS (ES+) m/z 657.3 (MH+)

Anal. calculated for $C_{28}H_{29}FN_4O_4S_2 \cdot 1.2C_6H_8O_7$ (Citric acid). $2.6H_2O$

	C	50.4	H	5.6	N	6.0
Found	C	50.1	H	5.4	N	6.3

5 **Example 44**

(2S)-2-{5-[1-(Thiazol-2-yl) -2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid

Treatment of tert-butyl (2S)-2-{5-[1-(thiazol-2-yl) -2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate with TFA by a similar method to that used for Example 39 gave the HCl salt of the title compound in 73 % yield.

1H NMR (DMSO- d_6 , 250MHz) δ : 2.0-2.3 (2H,m) ; 2.45-2.55 (partially obscured by the solvent signals, m); 2.55-2.9 (7H,m); 2.94 (3H,d); 3.1-3.28 (2H,m); 4.48-4.5 (1H,m); 4.82-4.94 (2H,m); 6.2-6.3 (1H,m) ; 6.88-7.02 (4H, m) ; 7.05-7.2 (3H,m) ; 7.62 (1H,t); 7.75 (2H,m); 7.86 (1H,d); 8.64-8.72 (1H,t); 9.16(1H,m).

15 MS (ES-) m/z 599.1 (MH-)

Anal. calculated for $C_{28}H_{29}FN_4O_6S_2 \cdot HCl \cdot 0.4H_2O \cdot 1.0C_6H_8O_7$ (Citric acid).

	C	45.3	H	5.1	N	6.2
Found	C	45.2	H	4.7	N	6.1

20 **Example 45**

Methyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate

The title compound was obtained from 2-(4-fluorophenethyl)-5-methoxybenzoic acid and L-methionone methyl ester in 54 % yield by a similar method to that used for Example 25 40.

1H NMR (DMSO- d_6 , 300MHz) δ : 1.87-2.08 (5H,m) ; 2.54 (3H,s); 2.58-2.9 (6H,m); 3.6 (3H,s); 4.44 -4.65 (3H,m) ; 5.76-5.84 (1H, m) ; 6.78-6.82 (2H,m) ; 6.95-7.06 (3H,m); 7.1-7.2 (4H,m); 7.4-7.58 (3H,m); 7.6 (1H,d); 8.72 (1H,t).

MS (ES+) m/z 608.4 (MH+)

30 The starting material was prepared as follows :

A mixture of 2-methylimidazole (8.0g, 97.5 mmol) and 4-fluorophenacyl bromide (21.16g, 97.5 mmol) dissolved in ethyl acetate (40ml) at ambient temperature was treated with triethylamine (16.4 ml, 117 mmol) and the cloudy mixture heated at reflux for 16 hours. The mixture was cooled to ambient temperature and diluted with ethyl acetate and water, washed
5 with water and brine, dried and evaporated to dryness. The residue was recrystallised (3x) from dichloromethane/methanol/isohexane to give 1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethanone (1, 9.0g, 42%).

¹H NMR (DMSO d₆, 300MHz) δ: 2.12 (3H,s); 5.64 (2H, s); 6.72 (1H,s); 6.96 (1H,s); 7.34 (2H,t); 8.12 (2H,dd).

10 MS (ES+) m/z 219 (MH)+.

A cooled (ice-bath) solution of 1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethanone (9.0g, 41.3 mmol) in methanol (60 ml) was treated portionwise with sodium borohydride (1.56g, 41.3 mmol) and the mixture stirred at ambient temperature for 16 hours. Most of the methanol was removed and water and ethyl acetate added. The organic layer was separated,
15 the aqueous extracted with ethyl acetate (2x) and the combined organic extracts washed with water, brine, dried filtered and evaporated to give a residue which was triturated with diethyl ether to give 1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethanol (7.7g, 84%).

¹H NMR (CDCl₃, 300MHz) δ 2.14 (3H,s); 3.98 (2H,d); 4.88 (1H,t); 6.74 (1H,s); 6.8 (1H,s); 7.04 (2H, t); 7.21- 7.32 (2H,m).

20 MS (ES+) m/z 221 (MH)+.

DEAD (1.95g, 12 mmol) added to a mixture of 1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethanol (2.48g, 11 mmol), methyl 2-(4-fluorophenethyl)-5-hydroxybenzoate (3.4g, 12 mmol), and triphenyl phosphine (3.3g, 12 mmol), in THF (60ml). The mixture was cooled in an ice-bath. After stirring for 16 hours the mixture was
25 evaporated to dryness and the residue purified by chromatography on silica (90b Biotage column, gradient elution 1-20% methanol/dichloromethane). Appropriate fractions were collected and evaporated to give methyl 5-[1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethoxy]-2-4-fluorophenethylbenzoate (4.2g, ~75%) which was used without further purification.

¹H NMR (CDCl₃, 300MHz) δ 2.21 (3H,s); 2.76 (2H, t); 3.08 (2H,t); 4.12-4.32 (1H,m); 5.3 (1H,t); 6.72-6.8 (1H, m); 6.84-6.98 (5H,m); 7.0-7.14 (4H,m); 7.16-7.24 (2H,m); 7.3 (1H,d).

MS (ES+) m/z 477 (MH)+; 279 (MH)+ for Ph₃ O.

- 5 Methyl 5-[1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoate (4.2g, ~8.8 mmol) was hydrolysed with aqueous sodium hydroxide solution under the usual conditions to give 5-[1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoic acid (1.84g, ~45%) as a colourless gum.

¹HNMR (DMSO d₆ , 300MHz) δ 2.52 (3H,s); 2.68 (2H,t); 3.0 (2H, dd); 4.48- 2.04 (2H,m) ; 5.76-5.86 (1H,m); 6.9-6.98(1H, m); 7.0-7.28(9H,m); 7.48 (2H,t); 7.62 (1H,d).

MS (ES+) m/z 463 (MH)+.

Example 46

(2S)-2-{5-[1-(4-Fluorophenyl)-2-(2-methylimidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyric acid

- 15 Treatment of methyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate with aqueous sodium hydroxide followed by the usual isolation procedure gave the hydrochloride salt of the title compound as a white solid in 64 % yield
- 20 ¹H NMR (DMSO-d₆, 300MHz) δ : 1.9-2.05 (5H,m) ; 2.57 (3H,s); 2.6-2.72 (2H,m); 2.75-2.85 (2H,m); 4.4-4.7 (3H,m); 5.78-5.86 (1H, m); 6.78-6.84 (2H,m) ; 6.95-7.08 (3H,m); 7.14-7.28 (4H,m); 7.45-7.55 (3H,m); 7.66 (1H,s); 8.58 (1H,t).
- MS (ES+) m/z 592 (MH+)

Example 47

tert-Butyl (2S)-2-{4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenyl)benzoylamino}-4-methylsulfanylbutyrate

- The title compound was prepared, in 33% yield, from 5-(1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy)-2-(4-fluorophenethyl)benzoic acid and L-methionine tert-butyl ester by a
- 30 similar method to that used for Example 38.

^1H NMR (DMSO- d_6 , HOAc- d_4 , 300MHz) δ : 1.34 (9H,s) ; 1.64-1.8 (2H,m); 1.92 (3H,d); 2.04-2.28 (2H,m); 2.6-2.74 (4H,dd); 4.08-4.18 (1H, m) ; 4.48 (2H,bds) ; 5.82 (1H,bds); 6.82-6.94 (2H,m); 7.04-7.2 (6H,m); 7.2-7.38 (4H,m); 7.4-7.52 (2H,m).

MS (ES+) m/z 608.3 (MH+)

5 Anal.calculated for $\text{C}_{33}\text{H}_{35}\text{F}_2\text{N}_3\text{O}_4\text{S} \cdot 0.9\text{C}_6\text{H}_8\text{O}_7$ (Citric acid).1.4 H_2O

	C	57.2	H	5.6	N	5.2
Found	C	57.1	H	5.2	N	5.3

The starting material was prepared as follows:

Sulphuryl chloride (4.4ml) was carefully added to a solution of 4-hydroxy-2-(4-fluorophenyl)benzoic acid (20.8g, 0.0896 mol) (from Example 32) in methanol (220 ml) and the mixture heated at reflux for 16 hours. The solvent was removed under reduced pressure and the residue partitioned between ethyl acetate and aqueous NaHCO_3 (pH ~8). The organic phase was washed with water, brine, dried (MgSO_4), filtered and evaporated to give a sticky brown solid which, on trituration with isohexane, gave methyl 4-hydroxy-2-(4-fluorophenyl)benzoate (14.34g, 65% yield).

^1H NMR (CDCl_3 , 300MHz) δ 3.64 (3H,s); 5.56 (1H,s); 6.74 (1H,d); 6.84 (1H,dd); 7.03-7.12 (2H,m); 7.16-7.28 (2H,m); 7.84 (1H,d).

MS (ES+) m/z 247 (MH)+.

Methyl 4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxy]-2-(4-fluorophenyl)benzoate was prepared from methyl 2-(4-fluorophenyl)-4-hydroxybenzoate and 2-(imidazol-1-yl)-1-(4-fluorophenyl)ethanol using the Mitsunobu reaction as described for the preparation of methyl 5-[2-(2-methylimidazol-1-yl)-1-(4-fluorophenyl)ethoxy]-2-(4-fluorophenethyl)benzoate.

^1H NMR (CDCl_3 , 300MHz) δ 3.6 (3H,s); 4.25-4.44 (2H,m); 5.4 (1H,t); 6.7- 6.78 (2H,m); 6.92 (1H,s); 7.0-7.3 (10H,m); 7.44 (1H,s); 7.76 (1H,d).

25 MS (ES+) m/z 435 (MH)+.

4-[2-(Imidazol-1-yl)-1-(4-fluorophenyl)ethoxy]-2-(4-fluorophenyl) benzoic acid was prepared from methyl 4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxy]-2-(4-fluorophenyl)benzoate by treatment with aqueous hydroxide in methanol and used without further purification.

30 ^1H NMR (DMSO d_6 , 300MHz) δ 4.7(2H,d); 6.02 (1H,t); 6.82-6.94 (2H,m); 7.1 -7.26(6H,m); 7.47(2H,dd); 7.62-7.68 (2H, m); 7.77 (1H,s); 9.16 (1H,s).

tert-Butyl (2S)-2-{4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-

The title compound was prepared, in 59 % yield, from 5-(1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy)-2-(4-fluorophenethyl)benzoic acid and L-methionine sulphone tert-butyl ester by a similar method to that used for Example 34.

MS (ES+) m/z 640.3 (MH+)

Found	C 56.1	H 5.60	N 5.80
-------	--------	--------	--------

Example 49

A solution of tert-butyl (2S)-2-{4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2(4-
20 fluorophenyl)benzoylamino}-4-methylsulfonylbutyrate (94mg, 0.15 mmol) in TFA (4ml) was stirred in an ice-bath for 3 hours. The TFA was evaporated under reduced pressure, the residue dissolved in the minimum amount of ethyl acetate / methanol , and 1M HCl/diethyl ether (5ml) added. The pale yellow precipitate obtained was triturated with iso-hexane, filtered, washed with iso-hexane and dried under vacuum to give the title compound (29mg).

MS (ES+) m/z 584.2 (MH+) for the acid, 598.2 (MH+) for the methyl ester.

Example 50**Methyl (2S)-2-{5-[2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate**

A mixture of 5-[2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoic acid (0.4g; 0.89mM), L-methionine methyl ester hydrochloride (0.27g, 1.35mM) and DMAP (0.54g, 4.43mM) was stirred until solution and EDC (0.026g, 1.35mM) in dichloromethane was added and the mixture stirred for 18 hours. It was purified on silica using dichloromethane then methanol/dichloromethane (5:95) as eluant to give the title product (0.47g, 89%) as a foam.

¹H NMR (CDCl₃, 300MHz) δ : 2.09 (3H,d) ; 2.15 (3H,d); 2.12 (1H,m); 2.25 (1H,m); 2.60 (2H,m) ; 2.81 (2H, m) ; 2.95 (2H,m) ; 3.68 (3H,d); 4.42 (1H,dq); 4.55 (1H,dt); 4.84 (1H,m); 5.71 (1H,m); 6.50 (1H,dd); 6.65 (1H,s); 6.82 (1H,q); 6.91 (3H,m); 7.07 (3H,m); 7.38 (2H,m), 7.85 (1H,d).

MS (ES+) m/z 597.4 (M+ 1 of C₃₀H₃₃FN₄O₄S₂ 596+1=597)

The starting material was prepared as follows :

A mixture of 4-methylimidazole (8.2g, 0.1M), anhydrous potassium carbonate (13.8g, 0.1M), potassium hydroxide (5.6g, 0.1M) and tetrabutylammonium chloride (1.13g, 3.51mM) in methylene chloride (170ml) at 0°C was treated with ethyl 2-bromoacetate (8.35g, 50mM). The mixture was stirred at 0°C for 1 hour and at ambient temperature, for 18 hours. It was then filtered and evaporated to dryness. The product was purified on silica using ethylacetate as eluant to give ethyl 4-methylimidazol-1-ylacetate and ethyl 5-methylimidazol-1-ylacetate (6.89g, 82%) as a 2:1 mixture.

¹H NMR (CDCl₃, 300 MHz) δ : 1.39 (t, 3H), 2.16 (s, 3H), 5-Me, 2.23 (s, 3H), 4-Me, 4.24 (2M, q), 4.58 (s, 2H), 5-Me isomer 4.60 (s, 2H), 4-Me isomer, 6.65 (s, 1H) 4-Me isomer, 6.80 (s, 1H), 5-Me isomer, 7.37 (s, 1H), 4-Me isomer, 7.42 (s, 1H), 5-Me isomer.

A solution of n-butyl lithium (1.6 M in hexanes) (21.7 ml, 34.72 mM) in dry diethylether (25 ml) was treated slowly with a solution of 2-bromothiazole (2.84 ml, 31.5 mM) in dry diethyl ether (50ml) at such a rate that the temperature did not exceed -65°C. A solution of ethyl 4-and 5-methylimidazol-1-ylacetate (6.89g, 41mM) in dry diethyl ether (25ml) was then added. The mixture was stirred at -70°C for 1½ hours, allowed to warm to ambient temperature over 3 hours and stirred for a further 18 hours. Saturated ammonium

chloride solution (120ml) was added slowly with stirring and the diethyl ether layer separated. The aqueous layer was extracted with diethyl ether and the combined organic extractions dried (MgSO_4) and evaporated to give a brown oil (5.85g). This was purified on silica using isohexane/ethyl acetate as eluant to give 2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethanone 5 (1.98g, 30%) as an oil which crystallised on standing.

^1H NMR (CDCl_3 , 300 MHz), δ : 2.26 (s, 3H), 5.49 (s, 2H), 6.69 (s, 1H), 7.44 (s, 1H), 7.80 (d, 1H), 8.07 (d, 1H).

MS m/z 208.0 ($\text{C}_9\text{H}_9\text{N}_3\text{OS}$ of M^{++1} requires 208)

A solution of 2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethanone (1.96g, 9.47mM) in 10 methanol (30ml) was cooled in an ice bath, treated with NaBH_4 (0.358g, 9.47mM) and stirred for 18 hours. Ethyl acetate (60ml) was then added, and the mixture washed with saturated aqueous NaHCO_3 solution, water, brine, dried (MgSO_4) and evaporated to give the product (0.56g). The aqueous layer was reextracted with ethyl acetate to give an additional product. The product was purified on silica eluting with methanol/dichloromethane (10:90) to give 2-15 (4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethanol (1.35g, 68%).

^1H NMR ($\text{DMSO}-d_6$, 300 MHz), δ : 2.03 (s, 3H), 3.15 (d, 1H), 4.08 (q, 1H), 4.34 (dd, 1H), 5.05 (m, 1H), 6.76 (s, 1H), 7.34 (s, 1H), 7.64 (t, 1H), 7.98 (t, 1H).

MS m/z 210.2 ($\text{C}_9\text{H}_{11}\text{N}_3\text{OS}$ of M^{++1} requires 210).

A stirred mixture of 2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethanol 20 (1.35g, 6.46 mM), methyl 2-(4-fluorophenethyl)-5-hydroxybenzoate (1.95g, 7.1 mM) and triphenylphosphine (1.86g, 7.1mM) in dry THF was cooled in an ice bath, and treated with DEAD (1.24g, 7.1mM). The mixture was allowed to warm to ambient temperature, stirred for 18 hours and evaporated to dryness. The product was purified on silica using methanol/dichloromethane (3 \rightarrow 10%) as eluant to give methyl 5-[2-(4-methylimidazol-1-yl)-25 1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoate (1.45g, 48%).

This was then dissolved in methanol (19ml) treated with 2M NaOH (9.4 ml), stirred for 18 hours, evaporated to dryness, redissolved in water (25ml) and concentrated HCl added dropwise until pH 6. The mixture was extracted with ethyl acetate (3 x 30ml), which was then washed with brine, dried (MgSO_4) and evaporated to dryness. The residue was purified 30 on silica using methanol/dichloromethane (10 \rightarrow 20%) to give 5-[2-(4-methylimidazol-1-yl)-

1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoate as a white foam on evaporating to dryness (0.96g, 68%).

¹H NMR (CDCl₃, 300 Mz), δ 2.21 (3H s), 2.85 (2H, t), 3.18 (2H, t), 4.44 (1H, q), 4.59 (1H, dd), 5.75 (1H, dd), 6.58 (1H, s), 6.80-6.94 (3H, m), 6.97 (1H, d), 7.12 (2H, dd), 7.37 (1H, d), 7.54 (1H, d), 7.85 (2H, d).

MS, m/z 450.3 (C₂₄H₂₂FN₃O₃S = 451 M⁺ requires 451).

Example 51

tert-Butyl (2S)-2-{5-[2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate

A mixture of 5-[2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoate (0.56g, 1.24mM), L-methionine tert-butyl ester (1.02g, 3.73mM), EDC (0.48g, 2.5mM), N-methylmorpholine (0.68ml, 7.31mM) and 1-hydroxybenzotriazole (0.25g, 1.85mM) was stirred in dry DMF (22ml) for 17 hours. It was then evaporated to dryness and the residue dissolved in dichloromethane, washed with saturated aqueous NaHCO₃ and purified on silica eluting with methanol/dichloromethane (5:95) to give the title compound as a white foam (0.68g, 82%).

¹H NMR (CDCl₃, 300 MHz) δ : 1.50 (9H, s), 2.14 (3H, d), 2.30 (1H, m), 2.52 (1H, m), 2.81 (2H, t), 2.92 (3H + 2H, s + m), 3.12 (1H, m), 3.27 (1H, m), 4.22 (1H, m), 4.55 (1H, dd), 5.72 (1H, m), 6.64 (2H, t), 6.80-6.94 (4H, m), 7.00-7.13 (3H, m), 7.37 (1H, t), 7.42 (1H, d), 7.84 (1H, d).

MS m/z 669.3 (C₃₃H₃₉FN₄O₆S₂ = 670 -1, M⁺ = 669)

Example 52

(2S)-2-{5-[2-(4-Methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid

A mixture of Example 50 (70 mg, 0.12mM), and 2M sodium hydroxide solution (0.7ml, 0.7mM) in methanol (2ml) was stirred for 18 hours. It was then evaporated to dryness and dissolved in diethyl ether. Hydrochloric acid (1M) was added to pH 1 and then excess NH₄OH (0.8%) was added to pH 10. The mixture was evaporated to dryness, extracted with dichloromethane, filtered and evaporated to give the title product as a foam (44.4mg, 65%).

¹H NMR (CDCl₃, 300MHz) δ :2.06 (3H, d), 2.13 (1H, m), 2.22 (3H, d), 2.33 (1H, m), 2.61 (2H, m), 2.72 (2H, m), 2.92 (2H, m), 4.51 (2H, m), 4.77 (1H, m), 5.74 (1H, m), 6.30 (1H, d), 6.73 (1H, qd), 6.86 (2H, t), 6.93-7.08 (5H, m), 7.37 (1H, m), 7.83 (1H, m), 7.92 (1H, d).

MS m/z 581.3 (C₂₉H₃₁FN₄O₄S₂ = 582 - 1 = 581 for M⁻)

5

Example 53

tert-Butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyrate

A mixture of L-methionine- tert-butyl ester (0.240 g), (8) (0.175 g.), DMAP (0.237 g.), EDC (0.149 g.) and DMF (10 ml.) was stirred under an inert atmosphere for 16 hrs. The DMF was evaporated off and the residue partitioned between 1M aqueous citric acid (10 ml) and dichloromethane (10 ml). The organic layer was separated, dried and applied directly to a silica flash column which was then eluted with ethyl acetate/methanol (9:1). The product was converted to the hydrochloride salt to give tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyrate as a white solid (0.145 g.).

¹H NMR(DMSO-d₆) δ 1.4(9H, s), 1.96(2H, m), 2.05(3H, s), 2.5(2H, m), 2.7(2H, m), 2.8 (2H, m), 3.3(2H, m), 3.8(3H, s), 4.4(1H, m), 5.7(1H, m), 6.8-7.66(12H, m), 8.6(1H, dd), 9.0(1H, s).

20 Anal. Calculated allowing for 1HCl, 0.5H₂O : C,62.2; H,6.2; N,6.0.

Found C,62.3; H,6.6; N,5.7.

MS(MH⁺) 650

Example 54

tert-Butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfonylbutyrate

tert-Butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfonylbutyrate was prepared using a similar method to that used for Example 53 except using L-methionine-sulphone-tert-butyl ester instead of L-methionine tert-butyl ester.

30

¹H NMR(DMSO-d₆) δ 1.39(9H, s), 2.16(2H, m), 2.7(2H, m), 2.85(2H, m), 3.0(3H, d), 3.25(4H, m), 3.8(3H, s), 4.4(1H, m), 5.7(1H, m), 6.82-7.55(12H, m), 8.68(1H, dd), 8.93(1H, s).

MS(MH⁺) 682

5 Anal. Calculated allowing for 1HCl, 1.5 H₂O : C,58.0; H,6.0; N,5.6.

Found C,58.2; H,6.0; N,5.4.

The starting material was prepared as follows:

A mixture of 1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethanone (1.3 g.), sodium borohydride (0.25 g.) and methanol (50 ml.) was stirred at ambient temperature for 2 hours
10 under an inert atmosphere. The methanol was evaporated away and water (30 ml.) added to the residue. The mixture was filtered and the solid washed with more water and dried to give 1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethanol (1.18 g.).

¹H NMR(DMSO-d₆) δ 2.84(2H, s), 3.4(3H, s), 4.75(1H, m), 5.43(1H, d), 6.56(1H, s), 7.2(2H, m), 7.32(2H, m), 7.48(1H, s).

15 MS(MH⁺) 221

DEAD (0.85 g.) was added over 10 minutes to a solution of 1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethanol (0.9 g.), methyl 2-(4-fluorophenethyl)-5-hydroxybenzoate and triphenylphosphine (1.3 g.) in tetrahydrofuran (30 ml.), cooled to 15°C under an inert atmosphere. The reaction was then stirred at ambient temperature for a further 16 hours. The
20 THF was evaporated away and the residue dissolved in dichloromethane and applied directly to a silica flash column which was then eluted with firstly ethyl acetate then ethyl acetate/methanol (9:1) to give 2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoate as a colourless gum (1.5 g.).

¹H NMR(CDCl₃) δ 2.78(2H, m), 3.1(3H, m), 3.23(1H, m), 3.4(3H, s), 3.85(3H, s), 5.29
25 (1H, t), 6.75-7.75(13H, m).

MS(MH⁺) 477.4

A mixture of 2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoate (1.5 g.), sodium hydroxide (0.62 g.), methanol (200 ml.) and water (30 ml.) was stirred at reflux for 16 hours. The mixture was then cooled, the methanol
30 evaporated away and the residue diluted with water (100 ml.). It was then washed with dichloromethane and the aqueous phase separated, acidified with 1M aqueous citric acid (25

ml.) and extracted with dichloromethane. The extracts were dried and evaporated to dryness to give 2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoic acid as a white foam (1.1 g).

5 Example 55

tert-Butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate and
tert-Butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate

10 The title compounds were synthesised from 2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoic acid using a similar method to that described in Example 53.

tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate :

¹H NMR(DMSO-d₆) δ 1.39(9H, s), 1.82(2H, m), 1.99(3H, s), 2.21(2H, m), 3.14(2H, m), 3.7(3H, s), 4.2(1H, m), 4.4(2H, s), 4.78(1H, m), 7.1-7.55(12H, m), 8.5(1H, d), 8.91(1H, s).

Anal. Calculated allowing for 1HCl, 1H₂O C,60.8; H,6.1; N,6.1.

Found C,60.8; H,5.8; N,5.9.

20 MS(MH⁺) 636.4

tert-butyl(2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate :

¹H NMR(DMSO-d₆) δ 1.4(9H, s), 2.0(1H, m), 2.1(1H, m), 2.7-3.5(4H, m), 2.92(3H, s), 3.72(3H, s), 4.28(1H, m), 4.41(2H, s), 4.8(1H, m), 7.1-7.58(12H, m), 8.7(1H, d), 8.97(1H, s), 14.35(1H, br.s).

Anal. Calculated allowing for 1HCl, 1.5H₂O C,57.4; H,5.7; N,5.7.

Found C,57.6; H,5.8; N,5.6.

MS(MH⁺) 668.3.

30 The starting material was prepared as follows:

Sodium hydride (0.089g., 60% dispersion in oil) was added to a suspension of 1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethanol (0.27 g.) in THF (50 ml.) and DMF (7 ml.) stirred under an inert atmosphere. The mixture was stirred for a further 30 minutes. A solution of methyl 4-bromomethyl-2-(4-fluorophenyl)benzoate (0.432 g.) in THF(20 ml.) was then added followed by tetrabutylammonium iodide (0.045 g.) and the mixture stirred for a further 30 minutes. It was then poured into ice/ saturated ammonium chloride solution and extracted with ethyl acetate. The extracts were dried, evaporated to dryness and the product purified by flash column chromatography eluting with firstly ethyl acetate, then ethyl acetate/methanol (9:1) to give 2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoate as a colourless gum (0.38 g.).

2-(4-Fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoate was hydrolysed to give 2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoic acid, using a similar method to that used to prepare 2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoic acid in Example 54.

Example 56

(2S)-2-{2-(4-Fluorophenyl)-6-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]pyrid-3-oylamino}-4-methylsulfanylbutyric acid

A solution of tert-butyl (2S)-2-{2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]pyrid-3-oylamino}-4-methylsulfanylbutyrate (0.1 g.) in TFA(20 ml.) was stirred at ambient temperature under an inert atmosphere for 30 minutes. The TFA was evaporated away and the residue was converted to the hydrochloride salt to give (2S)-2-{2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]pyrid-3-oylamino}-4-methylsulfanylbutyric acid as a white solid (0.092 g.).

¹H NMR(DMSO-d₆) δ 1.95(2H, m), 2.1(3H, d), 2.3(1H, m), 2.48(1H, m), 4.4(1H, m), 4.9(2H, m), 6.65(1H, m), 7.06(1H, d), 7.2-7.5(4H, m), 7.55-7.8(5H, m), 7.85(1H, d), 7.9(1H, m), 8.74(1H, dd), 9.29(1H, d).

Anal. Calculated allowing for 2HCl, 0.25Et₂O : C,54.0; H,4.7; N,8.7; S,5.0.

Found C,54.5; H,4.7; N,9.0; S,5.3.

MS(MH⁺) 553.3

The starting material was prepared as follows:

Thionyl chloride (3.8 g.) was added dropwise, over 10 minutes, to a solution of 2-chloropyridin-3-carboxylic acid HCl salt (1.0 g.) in methanol, stirred under an inert atmosphere and cooled in an ice/salt bath. The reaction was stirred for a further 5 days at ambient temperature. The methanol was evaporated away and the residue treated with saturated sodium bicarbonate solution and extracted with ethyl acetate. The extracts were dried and evaporated to dryness to give methyl 2-chloropyridin-3-carboxylate as a colourless liquid (0.9 g.)

$^1\text{H NMR}(\text{DMSO-}d_6) \delta$ 3.87(3H, s), 7.57(1H, dd), 8.25(1H, dd), 8.6(1H, dd).

10 MS(MH^+) 172

A mixture of methyl 2-chloropyridin-3-carboxylate (6.8 g.), 4-fluorobenzeneboronic acid (7.2 g.), 1M. aqueous sodium carbonate (40 ml.), $\text{Pd}(\text{dppb})\text{Cl}_2$ (1.2 g.), ethanol (10 ml.) and toluene (150 ml.) was stirred at reflux under an inert atmosphere for 5 hr. The reaction was then cooled, the organic layer separated, the aqueous extracted with toluene and then the combined extracts dried and evaporated to dryness. The product was purified by flash column chromatography eluting with ethyl acetate/iso-hexane (9:1, 8:2, 7:3) to give methyl 2-(4-fluorophenyl)pyridin-3-carboxylate as a white solid (7.5 g.)

$^1\text{H NMR}(\text{DMSO-}d_6) \delta$ 3.71(3H, s), 7.1(2H, t), 7.32(1H, dd), 7.52(2H, m), 8.1(1H, dd), 8.76(1H, dd).

20 MS(MH^+) 232.

A mixture of methyl 2-(4-fluorophenyl)pyridin-3-carboxylate (10.5 g.) and MCPBA (12.33 g., 65%) in dichloromethane was stirred at 0°C. for 1 hour and then at ambient temperature for 2 hours. It was then washed with saturated sodium bicarbonate solution (300 ml.), dried and applied directly to a silica flash column which was eluted firstly with ethyl acetate/iso-hexane (1:1), ethyl acetate and ethyl acetate/methanol (9:1) to give 2-(4-fluorophenyl)-3-(methoxycarbonyl)pyridine-1-oxide as a white solid (10.5 g.).

$^1\text{H NMR}(\text{CDCl}_3) \delta$: 3.63(3H, s), 7.18(2H, t), 7.31(1H, dd), 7.41(2H, t), 7.66(1H, d), 8.42(1H, d).

MS(MH^+) 248

30 A solution of 2-(4-fluorophenyl)-3-(methoxycarbonyl)pyridine-1-oxide (10.5 g.) in acetic anhydride (500 ml.) was stirred at reflux for 17 hr., cooled, evaporated to dryness and

the residue partitioned between saturated sodium bicarbonate solution and ethyl acetate. The organic layer was separated, dried and evaporated to dryness. The gum obtained was dissolved in methanol (400 ml.), treated with a solution of potassium carbonate (11.7 g.) in water (100ml) and stirred at ambient temperature for 2 hours. The methanol was evaporated off, the aqueous residue diluted with more water (200 ml.) and the mixture filtered to give methyl 2-(4-fluorophenyl)-6-hydroxypyridin-3-carboxylate (8.0 g.) as a brown solid.

¹H NMR(CDCl₃) δ 3.62(3H, s), 6.46(1H, d), 7.14(2H, t), 7.4(2H, dd), 7.98(1H, d), 10.93(1H, br.s).

MS(MH⁺) 248.2

10 Methyl 2-(4-fluorophenyl)-6-hydroxypyridin-3-carboxylate was reacted with 2-(imidazol-1-yl)-1-(4-fluorophenyl)ethanol by the procedure described for the preparation of 2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoate to give methyl 2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]pyridin-3-carboxylate.

15 ¹H NMR(CDCl₃) δ : 3.68(3H, s), 4.32(1H, dd), 4.44(1H, dd), 6.31(1H, t), 6.82(1H, d), 6.83(1H, s), 7.08(5H, m), 7.38(1H, s), 7.69(4H, m), 8.07(1H, d).

MS(MH⁺) 436.4.

Methyl 2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]pyridin-3-carboxylate was hydrolysed using sodium hydroxide to 2-(4-fluorophenyl)-6-[1-(4-

20 fluorophenyl)-2-(imidazol-1-yl)ethoxy]pyridin-3-carboxylic acid.

¹H NMR(DMSO-d₆) δ 4.48(1H, dd), 4.58(1H, dd), 6.37(1H, dd), 6.86(1H, s), 6.94(1H, d), 7.22(5H, m), 7.43(4H, m), 7.65(1H, s), 8.05(1H, d).

MS(MH⁺) 422.1

2-(4-Fluorophenyl)-6-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]pyridin-3-carboxylic acid was converted to tert-butyl (2S)-2-{2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]pyrid-3-ylamino}-4-methylsulfanylbutyrate by a similar method to that described for the preparation of tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyrate.

¹H NMR(DMSO-d₆) δ : 1.39(9H, d), 1.8(2H, m), 2.0(3H, d), 2.18(1H, m), 2.35(1H, m), 4.25(1H, m), 4.76(2H, m), 6.55(1H, m), 6.98(1H, d), 7.12-7.38(4H, m), 7.42-7.7(5H, m), 7.74(1H, d), 7.8(1H, m), 8.62(1H, dd), 9.15(1H, d).

MS(MH⁺) 609.3

Anal. Calculated allowing for : 2.0 HCl, 0.25 i.hexane : C,57.2; H,5.6; N,8.0; S,4.6.

Found C,57.5; H,5.5; N,8.2; S,4.9.

5 Example 57

tert-Butyl (2S)-2-{2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]pyrid-3-ylamino}-4-methylsulfanylbutyrate

- 2-(4-Fluorophenyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]pyrid-3-carboxylic acid (0.39g) and L-methionine-tert-butyl-ester.HCl (0.42g) were dissolved in DMF (50 ml) then DMAP (0.63g), EDC (0.25g) and HOBT (0.12g) were added under an inert atmosphere at ambient temperature. After 16 hours the solution was evaporated under reduced pressure, the residue obtained was diluted with 1M citric acid (10 ml) and extracted with 2% methanol/ dichloromethane (1x100ml, 1x60ml). The combined extracts were dried, filtered and concentrated under reduced pressure to give a yellow oil.
- 15 Purification by flash column chromatography eluting with methanol/ ethyl acetate (9:1) gave tert-butyl (2S)-2-{2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]pyrid-3-ylamino}-4-methylsulfanylbutyrate as a colourless foam (0.34g).
- ¹H NMR (CDCl₃) δ 1.43 (9H, s), 1.82-1.95 (2H, m), 2.04 (3H, s), 2.20-2.29 (2H, m), 2.88-2.99 (1H, m), 3.02-3.14 (1H, m), 3.20 (2H, s), 4.41-4.49 (1H, m), 4.55-4.66 (3H, m), 6.48 (1H, dd), 6.76 (1H, s), 7.00-7.15 (4H, m), 7.23-7.31 (4H, m), 7.58-7.66 (2H, m), 7.90 (1H, dd).
- Anal. Calculated allowing for 0.5 H₂O: C, 63.2 ; H, 6.1 ; N, 8.7;
Found C, 63.0 ; H, 5.8 ; N, 8.7 ;
MS (MH⁺) 637.4

The starting material was prepared as follows:

- 25 Methyl iodide (10.33 ml) was added to a stirred suspension of 2-chloro-6-nicotinic acid (22.8g) and potassium carbonate (36.8g) in DMF (190 ml), under an inert atmosphere at ambient temperature. After stirring vigorously for 16 hours the suspension was filtered and the solid residue washed with 10% methanol/ dichloromethane (100 ml). The filtrate was then concentrated under reduced pressure and the residue diluted with water (250 ml), extracted
30 with dichloromethane (3x150ml) and the combined extracts dried and concentrated under

reduced pressure. Purification by bulb to bulb distillation (140°C @ 0.05 mmHg) gave methyl 2-chloro-6-nicotinoate as a white solid (23.6g).

¹H NMR (CDCl₃) δ 2.50 (3H, s), 3.85 (3H, s), 7.10 (1H, d), 8.00 (1H, d).

MS(MH⁺) 186

5 Methyl 2-chloro-6-nicotinoate (23.1 g) and 4-fluorobenzeneboronic acid (22.6g) were dissolved in ethanol (25 ml) and toluene (350 ml) and Pd(dppb)Cl₂ (3.76g) were added under an inert atmosphere. An aqueous solution of 1M, sodium carbonate (124 ml) was added and the suspension heated at reflux for 4 hours and cooled to ambient temperature. The organic layer was separated and the aqueous layer extracted with dichloromethane (3x100 ml). The
10 combined organic extracts were dried, filtered and concentrated under reduced pressure to give a yellow oil. Purification by flash column chromatography eluting with ethyl acetate/ isohexane (4:6) gave methyl 2-(4-fluorophenyl)-6-methylpyridin-3-carboxylate as a pale yellow solid (27.8g).

¹H NMR (CDCl₃) δ 2.64 (3H, s), 3.68 (3H, s), 7.05-7.14 (2H, m), 7.18 (1H, d), 7.45-7.53
15 (2H, m), 8.03 (1H, d).

MS(MH⁺) 246

Methyl 2-(4-fluorophenyl)-6-methylpyridin-3-carboxylate (17.3g) was dissolved in dichloromethane (600 ml), the solution cooled to 0°C and MCPBA (24.3g) added in portions over 25 minutes. The solution was then warmed to ambient temperature and stirred for 16
20 hours washed with saturated aqueous sodium bicarbonate (500 ml), dried and concentrated under reduced pressure. The residue was purified by flash column chromatography, eluting with methanol/ ethyl acetate (1:12) to give 2-(4-fluorophenyl)-3-(methoxycarbonyl)-6-methylpyridine-1-oxide as a pale yellow oil which crystallised on standing (18.6g).

¹H NMR (CDCl₃) δ 2.58 (3H, s), 3.62 (3H, s), 7.11-7.18 (2H, m), 7.33-7.42 (3H, m), 7.63
25 (1H, d).

MS(MH⁺) 261

Methane sulfonyl chloride (7.11 ml) was added to a stirred solution of 2-(4-fluorophenyl)-3-(methoxycarbonyl)-6-methylpyridine-1-oxide (12.0g) in toluene (300 ml) and the mixture heated at reflux under an inert atmosphere for 16 hours. The solution was
30 partially concentrated under reduced pressure and then applied directly to a silica flash

column and eluted with ethyl acetate/ iso-hexane (1:5) to give methyl 2-(4-fluorophenyl)-6-chloromethylpyridin-3-carboxylate as a white crystalline solid (13.6g).

¹H NMR (CDCl₃) δ 3.72 (3H, s), 4.72 (2H, s), 7.08-7.16 (2H, m), 7.49-7.58 (3H, m), 8.14 (1H, d).

5 MS(MH⁺) 280

Sodium hydride (0.096g) was added to a stirred solution of 1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethanol (0.53g) in DMF (20 ml) under an inert atmosphere at ambient temperature. After 30 minutes the solution was cooled to -40°C and a solution of methyl 2-(4-fluorophenyl)-6-chloromethylpyridin-3-carboxylate (0.67g) in DMF (5 ml) added before
10 warming to 0°C over 90 minutes. Saturated aqueous ammonium chloride (1 ml) was added and the suspension concentrated under reduced pressure to give a yellow residue. Saturated brine (15 ml) was added and the mixture extracted with 3% methanol/ dichloromethane (5x25ml). The combined extracts were dried, and concentrated under reduced pressure.

Purification by flash column chromatography eluting with methanol/dichloromethane (1:12)

15 gave methyl 2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]pyrid-3-carboxylate as a yellow gum (0.30g).

¹H NMR (CDCl₃) δ : 2.95-3.17 (2H, m), 3.38 (3H, s), 3.71 (3H, s), 4.45-4.65 (3H, m), 6.86 (2H, s), 6.97-7.16 (4H, m), 7.21-7.33 (3H, m), 7.36 (1H, s), 7.40-7.50 (2H, m), 8.09 (1H, d).

MS(MH⁺) 464

20 A mixture of methyl 2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]pyrid-3-carboxylate (0.47g), sodium hydroxide (0.20g), water (6 ml) and methanol (30 ml) was heated at reflux for 3 hours and then cooled to ambient temperature. The mixture was concentrated under reduced pressure, 1M citric acid (6 ml) and water (6 ml) were added and the mixture extracted with 5% methanol/ dichloromethane (3x50ml). The
25 extracts were dried and concentrated under reduced pressure to give 2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]pyrid-3-carboxylic acid as a yellow gum (0.39g).

MS(MH⁺) 450

Example 58**tert-Butyl (2S)-2-{3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyrid-2-ylamino}-4-methylsulfanylbutyrate**

3-(4-Fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyridin-2-carboxylic acid (0.61g) and L-methionine-tert-butyl-ester.HCl (0.64g) were dissolved in dichloromethane (50 ml) then DMAP (0.96g) and EDC (0.38g) were added under an inert atmosphere at ambient temperature. After stirring for 16 hours the solution was washed with 1M citric acid (60 ml) and the organic layer dried and concentrated under reduced pressure. Purification on a silica flash column eluting with ethyl acetate gave tert-butyl (2S)-2-{3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyrid-2-ylamino}-4-methylsulfanylbutyrate as a colourless foam (0.27g).

¹H NMR (CDCl₃) δ : 1.54 (9H, d), 1.75-1.98 (2H, m), 2.05 (3H, s), 2.12-2.29 (1H, m), 2.33-2.63 (1H, m), 2.76-2.88 (2H, m), 3.06-3.30 (4H, m), 3.42 (3H, s), 4.63-4.73 (1H, m), 6.05-6.21 (1H, m), 6.76-7.03 (5H, m), 7.06-7.16 (2H, m), 7.26-7.42 (4H, m), 8.06-8.14 (1H, m).

Anal. Calculated allowing for 1.5 HCl : C, 59.6 ; H, 5.9 ; N, 7.9.

Found C, 59.8 ; H, 5.9 ; N, 7.7 ;

MS(MH⁺) 651

The starting material was prepared as follows:

Methyl 3-(trifluoromethanesulfonyloxy)pyridin-2-carboxylate was prepared in two steps from 2-hydroxypicolinic acid using the procedure of Subramanyam, C.; Chattarjee, S.; Mallamo, J.P. Tetrahedron Lett., 1996, 37, 459.

Methyl 3-hydroxypyridin-2-carboxylate:

¹H NMR (CDCl₃) δ : 4.08 (3H, s), 7.40 (2H, m), 8.28 (1H, dd).

MS(MH⁺) 154

Methyl 3-(trifluoromethanesulfonyloxy)pyridin-2-carboxylate:

¹H NMR (CDCl₃) δ : 4.07 (3H, s), 7.61 (1H, dd), 7.74 (1H, dd), 8.76 (1H, d).

MS(MH⁺) 286

Tributylamine (24.6 ml) was added to a stirred mixture of methyl 3-(trifluoromethanesulfonyloxy)pyridin-2-carboxylate (19.6g), bis (triortho-

tolyphosphine)palladium(II) chloride (1.08g), copper (I) iodide (0.13g) and 4-fluorostyrene (12.3 ml) in DMF (300 ml). The solution was heated under an inert atmosphere at 90°C for 26 hours then cooled to ambient temperature and concentrated under reduced pressure. The residue was dissolved in dichloromethane (300 ml), the solid filtered and the filtrate washed
5 with 1M. citric acid (3x100 ml), saturated aqueous sodium bicarbonate (150 ml), dried and concentrated under reduced pressure. The brown oil obtained was pre-absorbed onto silica then purified on a silica flash column eluting with ethyl acetate/ iso-hexane (4:6) to give methyl 3-[2-(4-fluorophenyl)ethenyl]pyridin-2-carboxylate as a pale yellow solid (12.4g).
¹H NMR (CDCl₃) δ : 4.02 (3H, s), 7.01 (1H, d), 7.04-7.11 (2H, m), 7.46 (1H, dd), 7.49-7.56
10 (2H, m), 7.83 (1H, d), 8.05 (1H, d), 8.59 (1H, d).
MS(MH⁺) 258

A suspension of 10% palladium on carbon (2.5g) and methyl 3-[2-(4-fluorophenyl)ethenyl]pyridin-2-carboxylate (12.4g) in ethyl acetate (200 ml) was stirred vigorously under an hydrogen atmosphere at ambient temperature for four hours. The
15 suspension was filtered through Celite and the filtrate concentrated under reduced pressure to give methyl 3-(4-fluorophenethyl)pyridin-2-carboxylate as a mobile, pale yellow oil (11.72g).
NMR data (CDCl₃) δ 2.89 (2H, dd), 3.21 (2H, dd), 3.97 (3H, s), 6.89-7.00 (2H, m), 7.06-7.16 (2H, m), 7.33 (1H, dd), 7.49 (1H, d), 8.58 (1H, dd).
MS(MH⁺) 260

20 Methyl 3-(4-fluorophenethyl)pyridin-2-carboxylate (11.7g) was dissolved in dichloromethane (600 ml), cooled to 0°C and MCPBA (14.7g) added in portions over 20 minutes. The solution was then warmed to ambient temperature and stirred for 16 hours. The mixture was washed with saturated aqueous sodium bicarbonate (2x200 ml), dried and then concentrated under reduced pressure. The residue was purified on a silica flash column eluting
25 with methanol/dichloromethane (1:12) to give 3-(4-fluorophenethyl)-2-(methoxycarbonyl)pyridine-1-oxide as a pale yellow oil which was crystallised from diethyl ether (10.5g).
¹H NMR (CDCl₃) δ 2.77-2.92 (4H, m), 4.03 (3H, s), 6.91-7.01 (3H, m), 7.03-7.09 (2H, m), 7.16 (1H, d), 8.14 (1H, d).
30 MS(MH⁺) 276

A solution of 3-(4-fluorophenethyl)-2-(methoxycarbonyl)pyridine-1-oxide (7.96g) in DMF (100 ml) and trifluoroacetic anhydride (40.9 ml) was stirred under an inert atmosphere at ambient temperature for 24 hours. The reaction mixture was concentrated under reduced pressure, diluted with saturated aqueous sodium bicarbonate (150 ml) and extracted with 5 dichloromethane (3x100ml). The combined organic extracts were dried and concentrated under reduced pressure and the residue purified on a silica flash column eluting with methanol/ dichloromethane (1:9) to give methyl 3-(4-fluorophenethyl)-6-hydroxypyridin-2-carboxylate as a cream solid (7.7g).

¹H NMR (CDCl₃) δ 2.80 (2H, dd), 3.08 (2H, dd), 3.95 (3H, s), 6.71 (1H, d), 6.92-7.01 (2H, 10 m), 7.06-7.11 (2H, m), 7.28 (1H, d), 9.63 (1H, br. s).

MS(MH⁺) 276

DEAD (0.85ml) was added dropwise over 2 minutes to a stirred suspension of methyl 3-(4-fluorophenethyl)-6-hydroxypyridin-2-carboxylate (1.0g), 2-(1-methylimidazol-5-yl)-1-(4-fluorophenyl)ethanol (0.80g) and triphenylphosphine (1.43g) in THF (70 ml) under an inert 15 atmosphere at ambient temperature. After stirring for 16 hours the solution was concentrated under reduced pressure and loaded directly onto a silica flash column eluting with methanol/ dichloromethane (1:20) to give methyl 3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyridin-2-carboxylate as a colourless oil (1.56g).

¹H NMR (CDCl₃) δ : 2.81 (2H, dd), 3.01-3.12 (3H, m), 3.35 (1H, dd), 3.91 (3H, s), 6.21 (1H, 20 t), 6.68-6.81 (2H, m), 6.87-7.20 (5H, m), 7.26-7.37 (3H, m), 7.40-7.45 (1H, m).

MS(MH⁺) 478

A mixture of methyl 3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyridin-2-carboxylate (1.56g), sodium hydroxide (0.65g), water (12 ml) and methanol (80 ml) was heated at reflux for 16 hours and then cooled to ambient 25 temperature. The mixture was concentrated under reduced pressure, 1M citric acid (17 ml) added and the mixture extracted with 10% methanol/dichloromethane (1x50ml, 2x30ml). The extracts were dried and concentrated under reduced pressure to give 3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyridin-2-carboxylic acid as a pale yellow foam (1.22g).

Example 59**Cyclopentyl (2S)-2-{3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyrid-2-ylamino}-4-methylsulfanylbutyrate**

The title compound was prepared from 3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyridin-2-carboxylic acid by a similar route to that described for the preparation of tert-butyl (2S)-2-{3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyrid-2-ylamino}-4-methylsulfanylbutyrate but using L-methionine cyclopentyl ester rather than L-methionine-tert-butyl ester.

¹H NMR (CDCl₃) δ : 1.58-1.81 (7H, m), 1.84-1.99 (3H, m), 2.03-2.16 (1H, m) overlapping 2.07 (3H, d), 2.17-2.43 (3H, m), 2.58 (1H, m), 2.81 (2H, m), 3.08-3.37 (4H, m), 3.39 (3H, s), 4.64-4.78 (1H, m), 5.28 (1H, m), 6.05-6.18 (1H, m), 6.76-7.03 (6H, m), 7.11 (2H, m), 7.34 (3H, m), 8.10 (1H, m).

Anal. Calculated allowing for 0.5 H₂O: C, 64.4 ; H, 6.1 ; N, 8.3.

Found C, 64.4 ; H, 6.1 ; N, 8.2 ;

MS(MH⁺) 663

The starting material was prepared as follows :

A mixture of N-tert-butoxycarbonyl L-methionine (30 g.), cyclopentanol (31 g.), DMAP (44 g.), EDC (30 g.) and dichloromethane (200 ml.) was stirred under an inert atmosphere for 16 hours. It was then washed with IM. aqueous citric acid (200 ml.), saturated sodium carbonate solution (100 ml.) and brine, dried and evaporated to dryness to give N-tert-butoxycarbonyl L-methionine cyclopentyl ester as a gum (36 g.).

¹H NMR(CDCl₃) δ : 1.44(9H, s), 1.52-2.0(9H, m), 2.0-2.2(1H, m), 2.1(3H, s), 2.54(2H, m), 4.34(1H, m), 5.1(1H, br.s), 5.2(1H, m).

MS(MH⁺) 318

A mixture of N-tert-butoxycarbonyl L-methionine cyclopentyl ester (53 g.), TFA (200 ml.), triethylsilane (39 g.) and dichloromethane (2 L.) was stirred at ambient temperature for 2 hours, evaporated to dryness and then converted to the hydrochloride salt to give L-methionine cyclopentyl ester as a white crystalline solid (40.5 g.).

¹H NMR(DMSO-d₆) δ : 1.47-1.95(8H, m), 1.95-2.2(2H, m), 2.05(3H, s), 2.45-2.75(2H, m), 4.0(1H, t), 5.18(1H, m), 8.7(3H, br.s).

MS(MH⁺) 218

Example 60**(2S)-2-{2-(4-Fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyric acid**

5 A mixture of tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyrate (0.050g, 0.078 mmol) in TFA was stirred at ambient temperature under a nitrogen atmosphere for 4 hours. The TFA was evaporated away and the residues basified with saturated sodium bicarbonate solution, reacidified with 2NHCl to pH6 and extracted with dichloromethane. The extracts were dried
10 and evaporated to dryness. The residue was dried under high vacuum to give the title compound as a white solid (0.040g, 89 %).

¹H NMR (CDCl₃ + DMSO- d₆) δ : 2.05-2.15 (5H, m), 2.15-2.20 (1H, m), 2.20-2.30 (1H, m), 2.59-2.68 (2H, m), 2.81-2.90 (2H, d), 2.95-3.03 (2H, d), 3.50-3.61 (2H, m), 3.86-3.95 (3H, s),
15 4.73-4.83 (1H, m), 5.90 -6.01 (1H, m), 6.83-6.96 (3H, m), 7.01- 7.17 (4H, m), 7.18- 7.28 (1H, m), 7.35 -7.47 (3H, m), 7.83 -7.87 (1H, s), 9.08-9.13 (1H, m)

Anal. Calculated allowing for 5.5 HCl : C, 44.47; H, 4.70; N, 7.15; S, 8.19

Found C, 44.40; H, 4.70; N, 7.10; S, 7.80

MS (MH⁺). 583.3

20 The starting material was prepared as follows :

A mixture of 2-thiazole carboxaldehyde (25g, 221 mmol), triethylorthoformate (46mls, 41g, 276mmol) and para- toluenesulphonic acid (1.52g, 9 mmol) in ethanol (200 ml) was stirred for 16 hours under a nitrogen atmosphere at ambient temperature. A further amount of para- toluenesulphonic acid (2g, 11 mmol) was added and the reaction stirred for
25 another 16 hours. The mixture was treated with sodium bicarbonate to pH 9, stirred for a further 5 minutes and then filtered and the filtrate evaporated to dryness to give 2-(1,1-diethoxymethyl)thiazole as a clear oil, which was used without further purification (41g, 99%).

¹H NMR (CDCl₃) δ : 1.28 (6H, t), 3.60-3.80 (4H, m), 5.74 (1H, s), 7.34 (1H, d), 7.80 (1H, d).
30 MS (MH⁺). 187.4

Boron trifluoride etherate (21g, 18 ml, 148 mmol) was added dropwise to a mixture of 2-(1,1-diethoxymethyl)thiazole (27.6g 148 mmol) and triethyl phosphite (24.5g, 25mls, 148 mmol) in dichloromethane (300 ml) and the mixture refluxed for 12 hours under a nitrogen atmosphere. The reaction was allowed to cool to ambient temperature, treated with water (100 ml) and stirred for a further 10 minutes. The organic phase was separated, dried, filtered and evaporated to dryness. Purification by flash column chromatography eluting with ethyl acetate and ethyl acetate/methanol (9: 1 and 4:1) gave diethyl ethoxy(1,3-thiazol-2-yl)methyl phosphonate as a brown oil, (7g, 30 %).

¹H NMR (CDCl₃) δ : 1.23 - 1.36 (9H, m), 3.65- 3.73 (2H, m), 4.07- 4.30 (4H, m), 5.05 (1H, d), 7.42 (1H, d), 7.80 (1H, d).

MS (MH⁺). 280.2

A solution of n- butyl lithium (1.6M in hexane) (30 ml, 47 mmol) was added dropwise over 5 minutes to a solution of diethyl ethoxy(1,3-thiazol-2-yl)methyl phosphonate (15.21g, 55 mmol) in tetrahydrofuran (400 ml) cooled to -78 °C under a nitrogen atmosphere and the mixture was then stirred for a further 10 minutes. A solution of 1-methyl-1H-imidazole-5-carboxaldehyde (4) (4g, 36 mmol) in tetrahydrofuran (20 ml) was then added over 10 minutes to the reaction which was then stirred for a further 20 minutes, after the addition was completed. The reaction was treated with water (50 ml), allowed to warm to ambient temperature, acidified with hydrochloric acid (10N, 50 ml) and refluxed for 18 hours under a nitrogen atmosphere. The reaction was cooled to ambient temperature, basified with saturated aqueous sodium bicarbonate to pH 8, the tetrahydrofuran evaporated and the aqueous residues extracted with ethyl acetate (2 x 250 ml). The combined organic extracts were washed with brine, dried and evaporated to dryness. Purification by column flash column chromatography eluting with ethyl acetate/methanol (9:1 and 4:1) gave 2-(1-methylimidazol-5-yl)-1-(thiazol-2-yl)ethanone as a yellow solid, (7.3g, 97%).

¹H NMR (CDCl₃) δ : 3.65 (3H, s), 4.48 (2H, s), 7.00 (1H, s), 7.45 (1H, s), 7.73 (1H, s), 8.05 (1H, s).

MS (MH⁺). 208

Sodium borohydride (0.56g, 14.9 mmol) was added to a solution of 2-(1-methylimidazol-5-yl)-1-(thiazol-2-yl)ethanone in methanol (150 ml) and the mixture stirred

under a nitrogen atmosphere at ambient temperature for 2 hours. The reaction was evaporated to dryness and the resulting gum was treated with water (15ml) and stirred for 30 minutes to give a solid precipitate which was filtered and dried to give 2-(1-methylimidazol-5-yl)-1-(thiazol-2-yl)ethanol (2.67g, 95%).

¹H NMR (CDCl₃) δ : 2.88 -2.99 (1H, q), 3.11 -3.20 (1H, dd), 3.28 (2H, s), 3.51 (3H, s), 4.96 - 5.04 (1H, m), 6.36 (1H, s), 6.59 (1H, s), 7.60 (1H, d), 7.73 (1H, s).

MS (MH⁺). 210.3

A mixture of methyl 2-(4-fluorophenethyl)-5-hydroxybenzoate (1.83g, 6.7 mmol), 2-(1-methylimidazol-5-yl)-1-(thiazol-2-yl)ethanol (1.40g, 6.68 mmol) and triphenylphosphine (2.10g, 8.0 mmol) in tetrahydrofuran (25 ml) was stirred at ambient temperature under a nitrogen atmosphere. DEAD (1.39g, 1.25 ml 8.0 mmol) was added dropwise to the solution. (The reaction was maintained at ambient temperature ± 2° C). The orange solution was then stirred at ambient temperature under a nitrogen atmosphere for 16 hours. The tetrahydrofuran solution was evaporated to dryness to give an oil. Purification by flash column

chromatography eluting with ethyl acetate, ethyl acetate/methanol (9:1 and 4:1) gave methyl 2-(4-fluorophenethyl)-5-[2-(1-methylimidazol-5-yl)-1-(thiazol-2-yl)ethoxy]benzoate (2.07g, 67%) as a clear oil.

¹H NMR (CDCl₃) δ : 2.76 (2H, t), 3.10 (2H, t), 3.30- 3.45 (2H, m), 3.60 (3H, s), 3.90 (3H, s), 5.70 -5.75 (1H, m), 6.90-7.50 (10 H, m), 7.77 (1H, d).

MS (MH⁺). 466.4

A mixture of methyl 2-(4-fluorophenethyl)-5-[2-(1-methylimidazol-5-yl)-1-(thiazol-2-yl)ethoxy]benzoate (2.07g, 4.4 mmol) and sodium hydroxide (0.89g, 22.23 mmol) in methanol (100 ml) and water (9 ml) was stirred under a nitrogen atmosphere at reflux for 18 hours, cooled to ambient temperature and the methanol evaporated off. The residue was treated with aqueous citric acid (1M, 40 ml) and extracted with dichloromethane. The organic extracts were dried and evaporated to dryness to give 2-(4-fluorophenethyl)-5-[2-(1-methylimidazol-5-yl)-1-(thiazol-2-yl)ethoxy]benzoic acid as a white foam, (1.48g, 75%).

¹H NMR (CDCl₃) δ : 2.83 (2H, t), 3.20 (2H, t), 3.27- 3.53 (2H, m), 3.56 (3H, s), 4.30 -4.60 (1H, br), 5.70 -5.80 (1H, m), 6.83 - 7.16 (7H, m), 7.30 -7.66 (3H, m), 7.80 (1H, d).

MS (MH⁺). 452.4

A mixture of 2-(4-fluorophenethyl)-5-[2-(1-methylimidazol-5-yl)-1-(thiazol-2-yl)ethoxy]benzoic acid (0.74g, 1.63 mmol), DMAP (1.0g, 8.2 mmol), L- methionine tert-butyl ester HCl (1.0g, 4.92 mmol), EDC (0.63g, 3.27 mmol) and HOBt (0.22g, 1.63 mmol) in DMF (50 ml) was stirred at ambient temperature under a nitrogen atmosphere for 16 hours.

5 The reaction was evaporated to dryness and washed with aqueous citric acid (1M, 20 ml) and extracted with dichloromethane (20 ml). The extracts were washed with saturated brine and dried and applied directly onto a silica flash column which was eluted with ethyl acetate/methanol (9:1 and 4:1) to give tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyrate. The product
10 was dissolved in ethyl acetate and treated with 1M ethereal HCl (10 ml). The resulting solid was isolated by centrifuging, further washing with diethyl ether and finally drying under high vacuum to give tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyrate as a white solid, (0.57g, 54 % Yield).

15 ¹H NMR (CDCl₃) δ : 1.45- 1.53 (9H, s), 1.99- 2.10 (1H, m), 2.08 (3H, s), 2.16 - 2.31 (1H, m), 2.51- 2.63 (2H, t), 2.78 -2.97 (2H, m), 2.92 - 3.00 (2H, m), 3.30- 3.4 (2H, m), 3.57 (3H, s), 4.66 - 4.78 (1H, m), 5.68 - 5.73 (1H, m), 6.47 -6.55 (1H, t), 6.77- 7.13 (8H, m), 7.40- 7.45 (2H, m), 7.80 (1H, d).

Anal. Calculated allowing for : 2 HCl 2H₂O C, 53.00; H, 6.07; N, 7.49; S, 8.58

20 Found C, 53.00; H, 6.20; N, 7.00; S, 8.70

MS (MH⁺). 439.4

Example 61

(2S)-2-{2-(4-Fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-

25 **yl)ethoxy]benzoylamino}-4-methylsulfonylbutyric acid**

(2S)-2-{2-(4-Fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfonylbutyric acid was prepared from tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfonylbutyrate by a similar method to that used for Example 60.

30 ¹H NMR (CDCl₃ + DMSO- d₆) δ : 2.48-2.56 (1H, m), 2.78 -2.99 (2H, m), 2.92-3.03 (4H, m), 3.25 -3.34 (1H, m), 3.51 - 3.60 (1H, m), 3.88-3.97 (3H, s), 5.93- 6.05 (1H, m), 6.05- 6.15 (1H,

m), 6.88-6.98 (3H, m), 7.03 -7.18 (3H, m), 7.25- 7.54 (3H, m), 7.68- 7.95 (2H, m), 8.93 -9.04 (1H, m).

Anal. Calculated allowing for 1 H₂O, 4 HCl : C, 44.73; H, 4.79; N, 7.20; S, 8.2

Found C, 44.90; H, 5.00; N, 6.80; S, 7.00

5 MS (MH⁺). 615.3

The starting material was prepared as follows :

A mixture of 2-(4-fluorophenethyl)-5-[2-(1-methylimidazol-5-yl)-1-(thiazol-2-yl)ethoxy]benzoic acid (0.74g, 1.63 mmol), DMAP (1.40g, 11.5 mmol), L- methionine sulphone tert-butyl ester HCl (1.34g, 4.92 mmol), EDC (0.63g, 3.27 mmol) and HOBT 10 (0.22g, 1.63 mmol) in DMF (50 ml) was stirred at ambient temperature under a nitrogen atmosphere for 16 hours. The reaction was evaporated to dryness and washed with aqueous citric acid (1M, 20 ml) and extracted with dichloromethane (20 ml). The extracts were washed with saturated brine, dried and evaporated to dryness. Purification by flash column chromatography eluting with dichloromethane/methanol (95:5, 9:1/85:15) gave a gum. This 15 was redissolved in ethyl acetate and treated with 1M ethereal HCl (10 ml). The resulting solid precipitate was isolated by centrifuging, further washing with diethyl ether and drying under high vacuum to give tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfonylbutyrate as a white solid, (0.66g, 60 % Yield).

20 ¹H NMR (CDCl₃) δ : 1.51(9H, s), 2.22-2.38 (1H, m), 2.46- 2.61 (1H, m), 2.77 - 2.88 (2H, m), 2.92 (3H, s), 2.92 -3.04 (2H, m), 3.08- 3.42 (6H, m), 3.58 (3H, s), 4.65- 4.77 (1H, m), 5.66 - 5.73 (1H, m), 6.77 - 7.13 (11H, m), 7.25 - 7.35 (3H, m), 7.42 -7.51 (1H, m), 7.76 -7.83 (1H, m).

Anal. Calculated allowing for 2 HCl : C, 53.29 ; H, 5.56 ; N, 7.53 ; S, 8.62

25 Found C, 53.00 ; H, 5.60 ; N, 8.10 ; S, 7.80

MS (MH⁺). 671.4

Example 62**tert-Butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate**

A mixture of 2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoic acid (0.47g, 1.06 mmol) (from Example 63), DMAP (0.62g, 5.05 mmol), L-methionine tert-butyl ester HCl (0.62g, 3.02 mmol), EDC (0.39g, 2.02 mmol) and HOBT (0.137g, 1.01 mmol) in DMF (25 ml) was stirred at ambient temperature under a nitrogen atmosphere for 16 hours. The reaction was evaporated to dryness and washed with aqueous citric acid (1M, 10 ml) and extracted with dichloromethane (20 ml). The extracts
 10 were washed with saturated brine, dried and filtered. Purification by flash column chromatography eluting with dichloromethane/ methanol (95:5, 9:1 and 85:15) gave a gum. This was dissolved in ethyl acetate and treated with 1M ethereal HCl (10 ml). The resulting solid precipitate was isolated by centrifuging, further washing with diethyl ether and drying under high vacuum to give tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-
 15 methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate as a white solid, (0.366g, 55 %).

¹H NMR (CDCl₃) δ : 1.51(9H, s), 1.92- 2.08 (1H, m), 2.15 -2.33 (1H, m), 2.59- 2.70 (2H, m), 2.85-2.98 (5H, m), 3.00-3.23 (4H, m), 3.46 (3H, d), 4.34 (1H, d), 4.52 (1H, dd), 4.70 -4.83
 20 (1H, m), 4.83-4.90 (1H, m), 6.77 (1H, d), 6.99- 7.08 (3H, m), 7.12- 7.21 (5H, m), 7.38 (1H, d), 7.80 (1H, d).

Anal. Calculated allowing for : 2.75 H₂O, 1 HCl C, 55.27; H, 6.48; N, 7.58; S, 8.68

Found C, 55.00; H, 6.70; N, 7.50; S, 9.00

MS (MH⁺). 653.4

25

Example 63**tert-Butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate**

A mixture of 2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoic acid (0.47g, 1.01 mmol), DMAP (0.86g, 7.07 mmol), L-methionine
 30 sulphone tert-butyl ester HCl (0.83g, 3.03 mmol), EDC (0.39g, 2.02 mmol) and HOBT

(0.138g, 1.01 mmol) in DMF (25 ml) was stirred at ambient temperature under a nitrogen atmosphere for 16 hours. The reaction was evaporated to dryness and washed with aqueous citric acid (1M, 10 ml) and extracted with dichloromethane (20 ml). The extracts were washed with saturated brine and dried. Purification by flash column chromatography eluting with 5 dichloromethane/methanol (95:5, 9:1 and 85:15) gave a gum. This was dissolved in ethyl acetate and treated with ethereal HCl (10 ml). The resulting solid precipitate was isolated by centrifuging, further washing with diethyl ether and drying under high vacuum to give tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate as a white solid, (0.506g, 10 73%).

¹H NMR (CDCl₃) δ : 1.51 (9H, s), 2.30 -2.65 (2H, m), 2.87 -3.43 (7H, m), 2.94 (3H, s), 3.45 (3H, s), 4.30 (1H, d), 4.65(1H, dd), 4.70 - 4.87 (2H, m), 6.73 (1H, d), 6.89 -7.02 (2H, m), 7.10 -7.20 (4H, m), 7.25 (2H, d), 7.34 -7.55 (2H, m), 7.83 (1H, d).

Anal. Calculated allowing for 2 H₂O, 1.5 HCl : C, 52.65; H, 6.04; N, 7.22; S, 8.27

15 Found C, 53.00; H, 6.30; N, 7.20; S, 8.30

MS (MH⁺). 685.4

The starting material was synthesised as follows :

Sodium hydride, 60% dispersion in mineral oil (0.44g, 11 mmol) was washed with isohexane and was suspended in DMF (25 ml). A solution of 2-(1-methylimidazol-5-yl)-1-20 (thiazol-2-yl)ethanol (2.10g, 10 mmol) in DMF (75 ml) was added dropwise and the mixture stirred under a nitrogen atmosphere at ambient temperature for 10 minutes. A solution of methyl 5-bromomethyl-2-(4-fluorophenethyl)benzoate (3.51g, 10 mmol) in DMF (50ml) was added dropwise to the reaction followed by tetrabutylammonium iodide (0.37g, 1 mmol) and the mixture was then stirred at ambient temperature for 16 hours. The reaction was evaporated 25 to dryness and the residue treated with water (100 ml) and extracted with dichloromethane (100 ml). The extracts were washed with saturated brine dried and evaporated to dryness. Purification by flash column chromatography eluting with dichloromethane/methanol (95:5 and 9:1) gave methyl 2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoate as a yellow gum, (3.32g, 69 %).

¹H NMR (CDCl₃) δ : 2.85 (2H, t), 3.12- 3.30 (4H, m), 3.50 (3H, s), 3.93 (3H, s), 4.45 (1H, d), 4.62 (1H, d), 4.90 - 4.97 (1H, m), 6.87 (1H, s), 6.95-7.05 (3H, m), 7.13 (3H, m), 7.40 (2H, d), 7.80 (2H, m).

MS (MH⁺). 480.4

5 A mixture of methyl 2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoate (3.32g, 6.92 mmol), sodium hydroxide (1.38g, 34.61 mmol) in methanol (150 ml) and water (15 ml) was stirred under a nitrogen atmosphere at reflux for 18 hours, cooled to ambient temperature and the methanol evaporated off. The residue was treated with aqueous citric acid (1M, 40 ml) and extracted with dichloromethane. The organic
10 extracts were dried and evaporated to dryness to give 2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoic acid as a yellow foam (2.85, 88%).

¹H NMR (CDCl₃) δ : 2.85- 2.98 (2H, m), 3.04 - 3.34 (4H, m), 3.53 (3H, s), 4.34 (1H, d), 4.72 (1H, d), 4.89 (1H, dd), 6.95 - 7.05 (2H, m), 7.04- 7.22 (4H, m), 7.33 -7.45 (2H, m), 7.57 (1H,
15 s), 7.83 (2H, d), 8.00 (1H, br).

MS (MH⁺). 466.4

Example 64

tert-Butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate
20

A mixture of 2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoic acid (0.44g, 1.07 mmol) (from Example 65), DMAP (0.62g, 5.03 mmol), L- methionine tert-butyl ester HCl (0.62g, 5.03 mmol), EDC (0.39g, 2.01 mmol) and HOBT (0.138g, 1.01 mmol) in DMF (25 ml) was stirred at ambient temperature under a
25 nitrogen atmosphere for 16 hours. The reaction was evaporated to dryness, the residue treated with aqueous citric acid (1M, 10 ml) and then extracted with dichloromethane (20 ml). The extracts were washed with saturated brine, dried and applied directly to a silica flash column eluting with dichloromethane/methanol (95:5, 9:1 and 85:15) to give a gum. This was dissolved in ethyl acetate and treated with ethereal 1M HCl (10 ml). The resulting solid was
30 isolated by centrifuging, further washing with diethyl ether and drying under high vacuum to

give tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate as a white solid, (0.373g, 59 %).

¹H NMR (DMSO- d₆) δ : 1.49 (9H, s), 1.80 -2.00 (2H, m), 2.06 (3H, s), 2.20 - 2.40 (2H, m), 3.13 -3.31 (4H, m), 4.22 - 4.32 (1H, m), 4.70 (2H, q), 5.11 (1H, q), 6.63 (1H, s), 7.20 -7.30 (2H, m), 7.30 -7.51(7H, m), 7.80 (1H, d), 7.80 (1H, d), 7.89 (1H, d), 8.53 (1H, d).

Anal. Calculated allowing for : 1.5 H₂O, 1.5 HCl C, 54.40; H, 5.92; N, 7.93; S, 9.08

Found C, 54.40; H, 6.00; N, 7.80; S, 9.00

MS (MH⁺). 625.4

10 Example 65

tert-Butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate

A mixture of 2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoic acid (0.44g, 1.07 mmol), DMAP (0.61g, 5.04 mmol), L-methionine
 15 tert-butyl ester HCl (0.71g, 3.03 mmol), EDC (0.39g, 2.01 mmol) and HOBT (0.138g, 1.01 mmol) in DMF (25 ml) was stirred at ambient temperature under a nitrogen atmosphere for 16 hours. The reaction was evaporated to dryness and washed with aqueous citric acid (1M, 10 ml) and extracted with dichloromethane (20 ml). The extracts were washed with saturated brine, dried and filtered. Purification by flash column chromatography eluting with
 20 dichloromethane/methanol (95:5, 9:1 and 85:15) gave a gum. This was dissolved in ethyl acetate and treated with ethereal 1M HCl (10 ml). The resulting solid was isolated by centrifuging, further washing with diethyl ether and drying under high vacuum to give tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate as a white solid, (0.406g, 62 %).

25 ¹H NMR (DMSO- d₆) δ : 1.40 (9H, s), 1.82 -2.02 (1H, m), 2.02-2.12 (1H, m), 2.73- 2.89 (1H, m), 2.93 (3H, s), 2.95 - 3.10 (2H, m), 3.10 - 3.25 (2H, m), 3.45 (3H, s), 4.18 4.28 (1H, m), 4.63 (2H, q), 5.02 (1H, q), 6.60 (1H, s), 7.10- 7.25 (2H, m), 7.25 -7.50 (7H, m), 7.71 (1H, d), 7.81 (1H, d), 8.79 (1H, d).

Anal. Calculated allowing for 1.5 HCl, 1.5 H₂O : C, 52.04; H, 5.66; N, 7.59; S, 8.68

30 Found C, 51.90; H, 5.40; N, 7.40; S, 8.70

MS (MH⁺). 657.4

The starting material was prepared as follows :

Sodium hydride 60% dispersion in mineral oil (0.17g, 4.34 mmol) was washed with isohexane and was suspended in DMF (25 ml). A solution of 2-(1-methylimidazol-5-yl)-1-(thiazol-2-yl)ethanol (0.83g, 3.94 mmol) in DMF (30 ml) was added dropwise and the mixture
5 stirred under a nitrogen atmosphere at ambient temperature for 10 minutes. A solution of methyl 4-bromomethyl-2-(4-fluorophenyl)benzoate (1.28 g, 3.94 mmol) in DMF (25mL) was added dropwise to the reaction followed by tetra butyl ammonium iodide (0.15g, 0.394 mmol) and the mixture was then stirred under a nitrogen atmosphere at ambient temperature for 16 hours. The reaction was evaporated to dryness, the residue treated with water (20ml) and
10 extracted with dichloromethane (50 ml). The extracts were washed with saturated brine, dried and filtered. Purification by flash column chromatography, eluting with dichloromethane/methanol (95:5 and 9:1) gave methyl 4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]-2-(4-fluorophenyl)benzoate (1.45g, 82 %) as a yellow gum.

15 ¹H NMR (CDCl₃) δ : 3.10 - 3.25 (2H, m), 3.49 (3H, s), 3.66 (3H, s), 4.50 (1H, d), 4.68 (1H, d), 4.98 (1H, q), 6.86 (1H, s), 7.03 - 7.13 (2H, m), 7.17 - 7.30 (4H, m), 7.32 (1H, s), 7.47 (1H, d), 7.78 (1H, d), 7.78 - 7.83 (1H, m).

MS (MH⁺). 451


A mixture of methyl 4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]-2-(4-fluorophenyl)benzoate (1.45g, 3.21 mmol) and sodium hydroxide (0.64g, 16.05 mmol) in
20 methanol (75 ml) and water (7 ml) was stirred under a nitrogen atmosphere at reflux for 18 hours, cooled to ambient temperature and the methanol evaporated off. The residue was treated with aqueous citric acid (1M, 5 ml) and extracted with dichloromethane. The organic extracts were dried and evaporated to dryness to give 2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-
25 (1-methylimidazol-5-yl)ethoxymethyl]benzoic acid as a white foam, (0.877g, 63%).

¹H NMR (CDCl₃) δ : 2.94 - 3.11 (2H, m), 3.43 (3H, s), 4.33 (1H, d), 4.66 (1H, d), 4.84 (1H, dd), 6.72 (1H, s), 6.89 - 7.13 (5H, m), 7.22 - 7.32 (3H, m), 7.55 (1H, s), 7.70 - 7.76 (2H, m).

MS (MH⁺). 438.4.


CLAIMS

1. A compound of Formula (1):




Formula (1)

wherein Ar¹ represents:




(A)

or



(B)

or



(C)

5

R⁵ is hydrogen, C₁₋₄alkyl, phenylC₁₋₄alkyl;

R⁶ is hydrogen, C₁₋₄alkyl, hydroxyC₁₋₄alkyl, haloC₁₋₄alkyl, dihaloC₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, sulfanylC₁₋₄alkyl, aminoC₁₋₄alkyl, N-(C₁₋₄alkyl)aminoC₁₋₄alkyl, N,N-di(C₁₋₄alkyl)aminoC₁₋₄alkyl or phenylC₁₋₄alkyl; m is 0, 1 or 2;

- 10 R¹² and R¹³ are independently hydrogen or C₁₋₄ alkyl;


Ar² is phenyl or heteroaryl;

p is 0 or 1;

Ar³ is phenyl, pyridinyl, pyridazinyl, pyrimidyl or pyrazinyl, the ring being substituted on ring carbon atoms by R² and -(CH₂)_nR³ and wherein Ar³ is attached to


- 15 Ar¹C(R¹²)R¹³CH(Ar²)O- by a ring carbon atom;

R² is a group of the Formula (2):



Formula (2)

wherein R^7 is hydrogen or C_{1-4} alkyl, R^8 is $-(CH_2)_q-R^{10}$ wherein q is 0-4 and R^{10} is C_{1-4} alkylsulfanyl, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, hydroxy, C_{1-4} alkoxy, carbamoyl, \underline{N} - C_{1-4} alkyl carbamoyl, $\underline{N,N}$ -(di C_{1-4} alkyl)carbamoyl, C_{1-4} alkyl, phenyl, thienyl, or C_{1-4} alkanoylamino, R^9 is hydroxy, C_{1-6} alkoxy, C_{3-9} cycloalkyloxy, heterocyclyloxy, 5 heterocyclyl C_{1-4} alkoxy or $-NH-SO_2-R^{11}$ wherein R^{11} represents, trifluoromethyl, C_{1-4} alkyl, phenyl, heteroaryl, aryl C_{1-4} alkyl or heteroaryl C_{1-4} alkyl; or R^2 represents a lactone of Formula (3):




Formula (3)

the group of Formula (2) or (3) having \underline{L} or \underline{D} configuration at the chiral alpha carbon in the 10 corresponding free amino acid; n is 0, 1 or 2; R^3 is phenyl or heteroaryl; phenyl and heteroaryl rings in R^3 , R^5 , R^6 , R^9 , R^{11} and Ar^2 are independently optionally substituted on ring carbon atoms in by up to three substituents selected from C_{1-4} alkyl, 15 halogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, amino, C_{1-4} alkylamino, di(C_{1-4} alkyl)amino, C_{1-4} alkanoylamino, nitro, cyano, carboxy, thiol, C_{1-4} alkylsulfanyl, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkanesulphonamido, \underline{N} -(C_{1-4} alkylsulphonyl)- \underline{N} - C_{1-4} alkylamino, aminosulfonyl, \underline{N} -(C_{1-4} alkyl)aminosulfonyl, $\underline{N,N}$ -di(C_{1-4} alkyl)aminosulfonyl, carbamoyl, \underline{N} -(C_{1-4} alkyl)carbamoyl, $\underline{N,N}$ - 20 (di C_{1-4} alkyl)carbamoyl, carbamoyl C_{1-4} alkyl, \underline{N} -(C_{1-4} alkyl)carbamoyl C_{1-4} alkyl, $\underline{N,N}$ -(di C_{1-4} alkyl)carbamoyl C_{1-4} alkyl, hydroxy C_{1-4} alkyl and C_{1-4} alkoxy C_{1-4} alkyl and on ring NH groups (replacing hydrogen) by C_{1-4} alkyl, C_{1-4} alkanoyl, C_{1-4} alkylsulfonyl, halo C_{1-4} alkyl, difluoromethyl or trifluoromethyl; or a pharmaceutically-acceptable salt, prodrug or solvate thereof.

25

2. A compound according to claim 1 wherein Ar^1 is of the formula (A) or (B):



wherein R^5 is hydrogen or methyl; R^6 is hydrogen, C_{1-4} alkyl, hydroxy C_{1-4} alkyl, amino C_{1-4} alkyl, fluoro C_{1-4} alkyl, difluoro C_{1-4} alkyl, C_{1-4} alkoxy or C_{1-4} alkoxy C_{1-4} alkyl and m is as defined in claim 1.

3. A compound according to either claim 1 or claim 2 wherein Ar^2 is phenyl, thiazolyl, pyridyl, triazolyl, pyrimidyl, pyrazinyl or pyridazinyl optionally substituted on ring carbon atoms by C_{1-4} alkyl, halo, nitro, cyano or C_{1-4} alkoxy C_{1-4} alkyl.


10

4. A compound according to either claim 1 or claim 2 wherein Ar^2 is phenyl or thiazolyl optionally substituted on ring carbon atoms by C_{1-4} alkyl, halo, nitro, cyano or C_{1-4} alkoxy C_{1-4} alkyl.

15 5. A compound according to any one of claims 1 to 4 wherein Ar^3 is phenyl or pyridyl, the ring being substituted on ring carbon atoms by R^2 and $-(CH_2)_nR^3$ and wherein Ar^3 is attached to $Ar^1C(R^{12})R^{13}CH(Ar^2)O-$ by a ring carbon atom.

6. A compound according to claim 5 wherein, when n is 0, the phenyl or pyridyl ring is substituted by R^2 in the 4-position and $-(CH_2)_nR^3$ in the 3- or 5-position and when n is 1 or 2, the phenyl or pyridyl ring is substituted by R^2 in the 3- or 5-position and $-(CH_2)_nR^3$ in the 4-position and wherein the positions indicated are relative to the point of attachment of Ar^3 to $-(CH_2)_p-$.

25 7. A compound according to any one of claims 1 to 6 wherein R^2 is of the formula (2):



wherein R⁷ is hydrogen or methyl;

R⁸ -(CH₂)_q-R¹⁰ wherein q is 1 or 2; R¹⁰ is C₁₋₄alkylsulfanyl, C₁₋₄alkylsulfinyl, C₁₋₄alkylsulfonyl, hydroxy or C₁₋₄alkoxy;

- 5 R⁹ is hydroxy, C₁₋₄alkoxy, C₃₋₉cycloalkyloxy, heterocyclyloxy or heterocyclylC₁₋₄alkoxy or of the formula -NHSO₂R¹¹ wherein R¹¹ is phenyl; wherein NH groups in heterocyclic groups in R⁹ are optionally substituted by methyl, ethyl, acetyl, propionyl, fluoromethyl, difluoromethyl or trifluoromethyl and ring carbon atoms in phenyl or heteroaryl groups in R¹¹ are optionally substituted by methyl, halo, C₁₋₄alkanoyl, nitro, cyano, C₁₋₄alkylsulfinyl, C₁₋₄alkylsulfonyl, carbamoyl, C₁₋₄alkylcarbamoyl or diC₁₋₄alkylcarbamoyl;
- 10 or wherein R² is of the formula (3) as defined in claim 1.

8. A compound according to any one of claims 1 to 7 wherein R³ is phenyl, pyridyl or thiazolyl and ring carbon atoms in R³ are optionally substituted by C₁₋₄alkyl, halo, C₁₋₄alkoxy, nitro, cyano or C₁₋₄alkoxyC₁₋₄alkyl and a ring NH group in a heteroaryl group in R³ is optionally substituted by C₁₋₄alkyl.
- 15

9. A compound according to any one of claims 1 to 8 wherein R¹¹ and R¹² are independently hydrogen or methyl.

20

10. A compound according to claim 1 which is:
- methyl (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
- (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoylamino]-4-methylsulfanylbutyric acid;
- 25 tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
- methyl (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(thiazol-2-yl)ethoxymethyl]benzoylamino}-4-methylsulfanyl butyric acid;

- (2S)-2-{2-(4-fluorophenyl)-4-[2-(imidazol-1-yl)-1-(thiazol-2-yl)ethoxymethyl]benzoylamino}-4-methylsulfanyl butyric acid;
methyl (2S)-2-{2-(4-fluorophenethyl)-5-[2-(imidazol-1-yl)-1-(4-fluorophenyl)-ethoxymethyl]benzoylamino}-4-methylsulfanyl butyrate;
- 5 (2S)-2-{2-(4-fluorophenethyl)-5-[2-(imidazol-1-yl)-1-(4-fluorophenyl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid;
methyl (2S)-2-{2-(4-(fluorobenzyl)-5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
(2S)-2-{2-(4-fluorobenzyl)-5-[1-(4-fluorophenyl)-2-(imidazol-1-
- 10 yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid;
methyl (2S)-2-{2-phenyl-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate; or
(2S)-2-{2-phenyl-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyric acid;
- 15 methyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-2-methyl-4-methylsulfanylbutyrate;
(2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-2-methyl-4-methylsulfanylbutyric acid;
N-(4-chlorobenzenesulfonyl)-(2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(imidazol-
- 20 1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyramide;
2-(morpholinomethyl)prop-2-yl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
methyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate;
- 25 (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyric acid;
tert-butyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate;
cyclopentyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-
- 30 fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate;
tert-butyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;


- 2-{5-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid;
- methyl (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate;
- 5 (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyric acid;
- tert-butyl (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate;
- tert-butyl (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
- 10 (2S)-2-{5-[1-(thiazol-2-yl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyric acid;
- methyl (2S)-2-{5-[1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate;
- 15 (2S)-2-{5-[1-(4-fluorophenyl)-2-(2-methylimidazol-1-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyric acid;
- tert-butyl (2S)-2-{4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenyl)benzoylamino}-4-methylsulfanylbutyrate;
- tert-butyl (2S)-2-{4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenyl)benzoylamino}-4-methylsulfonylbutyrate;
- 20 {4-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]-2-(4-fluorophenyl)benzoylamino}-4-methylsulfonylbutyric acid;
- methyl (2S)-2-{5-[2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyrate;
- 25 tert-butyl (2S)-2-{5-[2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfonylbutyrate;
- (2S)-2-{5-[2-(4-methylimidazol-1-yl)-1-(thiazol-2-yl)ethoxy]-2-(4-fluorophenethyl)benzoylamino}-4-methylsulfanylbutyric acid;
- tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyrate;
- 30 tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfonylbutyrate;

- tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
- tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate;
- 5 (2S)-2-{2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(imidazol-1-yl)ethoxy]pyrid-3-oylamino}-4-methylsulfanylbutyric acid;
- tert-butyl (2S)-2-{2-(4-fluorophenyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxymethyl]pyrid-3-oylamino}-4-methylsulfanylbutyrate;
- tert-butyl (2S)-2-{3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyrid-2-oylamino}-4-methylsulfanylbutyrate;
- 10 cyclopentyl (2S)-2-{3-(4-fluorophenethyl)-6-[1-(4-fluorophenyl)-2-(1-methylimidazol-5-yl)ethoxy]pyrid-2-oylamino}-4-methylsulfanylbutyrate;
- (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfanylbutyric acid;
- 15 (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxy]benzoylamino}-4-methylsulfonylbutyric acid;
- tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate;
- tert-butyl (2S)-2-{2-(4-fluorophenethyl)-5-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate;
- 20 tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfanylbutyrate; or
- tert-butyl (2S)-2-{2-(4-fluorophenyl)-4-[1-(thiazol-2-yl)-2-(1-methylimidazol-5-yl)ethoxymethyl]benzoylamino}-4-methylsulfonylbutyrate;
- 25 or a pharmaceutically-acceptable salt thereof.

11. A pharmaceutical composition which comprises a compound according to any one of claims 1 to 10 and a pharmaceutically-acceptable carrier.
- 30 12. A method of treating a disease or medical condition mediated through farnesylation of ras which comprises administering to a warm-blooded animal an effective amount of a compound according to any one of claims 1 to 10.

13. A compound according to any one of claims 1 to 10 for use as a medicament.
14. A compound according to any one of claims 1 to 10 for use in the preparation of a
5 medicament for treatment of a disease mediated through farnesylation of mutant ras.
15. A process for preparing a compound of the Formula I or a pharmaceutically acceptable salt, prodrug or solvate thereof, which process comprises:
deprotecting a compound of the formula (4):

10




(4)


wherein Ar^{1'} is Ar¹ as defined in claim 1 or protected Ar¹, Ar^{2'} is Ar² as defined in claim 1 or protected Ar² and Ar^{3'} is Ar³ as defined in claim 1 or protected Ar³; wherein at least one protecting group is present; and thereafter if necessary:

- 15 (i) forming a pharmaceutically-acceptable salt,
(ii) forming a prodrug, and/or
(iii) forming a solvate.

16. A process for preparing a compound of the Formula I or a pharmaceutically-acceptable
20 salt, prodrug or solvate thereof, which process comprises:
(i) reacting a compound of the formula (5) with a compound of the formula (6):




(5)



- 25 or (iii) converting one value of R⁹ in R² into another value of R⁹:

or (iii) reacting a compound in which R² in Ar^{3'} is carboxy with a compound of the formula (7):



wherein p, R⁷ and R⁸ are as defined in claim 1, Ar^{1'}, Ar^{2'}, Ar^{3'}, R⁷ and R⁸ are as defined in
5 claim 15, R²¹ is R⁹ as defined in claim 1 or a carboxy protecting group and when p is 1, L is a leaving group, and when p is 0, L is hydroxy; and thereafter if necessary:

- (i) removing any protecting groups;
- (ii) forming a pharmaceutically-acceptable salt, prodrug or solvate thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/03117

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D233/60 C07D233/64 C07D417/06 C07D401/12 A61K31/415

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 17070 A (UNIVERSITY OF PITTSBURGH) 15 May 1997 * the whole document, particularly example 179 *	1-16
A	WO 96 30015 A (MERCK & CO., INC.) 3 October 1996 * the whole document, particularly claims 12-15 *	1-16

☐ Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 January 1999

Date of mailing of the international search report

18/01/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Allard, M

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 98/03117

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 12
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 12 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 98/03117

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9717070 A	15-05-1997	AU 7597596 A EP 0873123 A	29-05-1997 28-10-1998
WO 9630015 A	03-10-1996	US 5578629 A US 5534537 A AU 5370196 A CA 2216564 A EP 0817630 A	26-11-1996 09-07-1996 16-10-1996 03-10-1996 14-01-1998